

Salinity problems in dredged sediments treated by phytoremediation

Floriane Sordes, Isabelle Techer, Slimane Sahli, Solène Cothelet, Allyson Moureaux, Chrystelle Bancon-Montigny, Jean-Claude Souche

▶ To cite this version:

Floriane Sordes, Isabelle Techer, Slimane Sahli, Solène Cothelet, Allyson Moureaux, et al.. Salinity problems in dredged sediments treated by phytoremediation. International symposium of LabEx DRIIHM, Sep 2021, Toulouse, France. , 2021, 10.34972/driihm-9b297b . hal-03346815

HAL Id: hal-03346815

https://hal.science/hal-03346815

Submitted on 21 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Salinity problems in dredged sediments

treated by phytoremediation


F. Sordes (1)*, I. Techer (1), S. Sahli (1), S. Cothelet (1), A. Moureaux (1), C. Bancon-Montigny (2), J.C. Souche (3) (1) Université de Nîmes, UPR CHROME, rue du Dr Georges Salan, 30021 Nîmes, France,











Metallic and organic contamination

Occitanie plan to dredge **5.8 millions m3** of sediments

26% of french dredged sediments are brought back

-> Chloride tend to accumulate in fine granulometry

inland and therefore are considered as waste.

64% of these sediments are silt or clay.

in the next 10 years.


(Hu et al. 2013)

Sediments and contaminants accumulate in the harbor. They both come from:

- Geological background Harbor activities
- Waste water Agriculture
- Industry



Drainage




Dredged sediments are naturally rich in different salts (Sea water chloride mean concentration: |Cl| = 20g/L).



Decrease of soluble contamination


Sediments drainage are usually observed (various drainage methods). After this chloride sediments concentration usually qualify them as non hazardous and non inert waste french category ([Cl] = 0.8-15 g/kg).




- concentration to an inert waste level in a reasonable amount of time ([Cl] < 0.8 mg/kg).
- Energetic valorisation of the plants by-products.

- Sediments need to be **dredged** to maintain touristic and economic activies.
- Level of heavy metals, PBC, HAP and TBT determine if dredged sediments should be brought back inland, instead of released offshore.
- Dredged sediments management face environmental, juridic and economic issues. Nearly one quarter of them are brought to land where they must be treated as waste. Regulation calls for a strict limitation of the offshore dumping.








Material and method


• 3 sites were studied : Pérols (65%<63µm, 5 months drainage), Port Camargue (58%<63µm, 2 month drainage) and le Grau du Roi (70%<63µm, totally drained).

• Arundo Donax plants were selected

Drainage in geotubes - Pérols (Sordes 2020)

Preliminar Survival test for Arundo donax in similar sediments regarding the granulometry. Sediments comes from Le Grau du Roi ([Cl] = 2.4 g/kg): Repoting vs planting from cutting.





- 2.5L repot into 7L of sediments
- Grow box (10 hour of light per day, 20°C, 0.5L water/pot/week)
- Harvest after 2 weeks and 4 months

Results Dryness 20cm under the surface (%) Chloride concentration (DW) drainage 15000 10000 Port Camargue Pérols Port Camargue Pérols

Occitanie dredging projects

Sediments are in hazardous waste category ([Cl] > 15 g/kg) which make it difficult to eliminate.

Occitanie Region (CC BY-SA 4.0)

In the long run, when facing high chloride level, Arbuscular Mycorrhiza modality shows the best results in bioconcentration and translocation in Arundo donax.

100% survival 0% survival

Results in accord with Pollastri et al. 2018 findings on salt stress reduction by Mycorrhyza.





Table 1: Comparison of above ground Cl accumulation in litterature		
Species	Cl accumulation	Authors
Suaeda Maritima	70 g/kg DW	Hasanuzzaman et al. 2014
Aeluropus littoralis	20 g/kg DW	Hasanuzzaman et al. 2014
Phragmites australis	24 g/kg DW	McSorley et al. 2016
Juncus maritimus	20 g/kg DW	Al Hassan et al. 2016
Typha latifolia	24 to 63 g/kg DW	Morteau et al. 2009; Rozema et al. 2014
Arundo donax	10 g/kg DW	This study

Conclusion

- Longer drainage should improve chloride elimination. Phyroremediation seems a good complementary technique.
- Tests realised with Arundo donax commonly used for energetic valorisation show good results in terms of uptake of chloride. Even with smaller bioaccumulation than other species, this specie have more interesting biomass production and economic valorization.
- Repot and Arbuscular Mycorrhiza show the best result in term of Cl- extraction and bioaccumulation.



References

- Al Hassan, M., del Pilar Lopez-Gresa, M., Boscaiu, M., & Vicente, O. (2016). Stress tolerance mechanisms in Juncus: responses to salinity and drought in three Juncus species adapted to different natural environments. Functional Plant Biology, 43(10), 949-960.
- Hasanuzzaman, M., Nahar, K., Alam, M., Bhowmik, P. C., Hossain, M., Rahman, M. M., ... & Fujita, M. (2014). Potential use of halophytes to remediate saline soils. BioMed research international, 2014.
- Hu, Y., Zhang, C., Wang, D. Z., Wen, J. Y., Chen, M. H., & Li, Y. (2013). Chloride ion transport and fate in oilfield wastewater reuse by interval dynamic multimedia aquivalence model. Water science and technology, 67(3), 628-634.
- McSorley, K., Rutter, A., Cumming, R., & Zeeb, B. A. (2016). Phytoextraction of chloride from a cement kiln dust (CKD) contaminated landfill with Phragmites australis. Waste management, 51, 111-118.
- Morteau, B., Triffault-Bouchet, G., Galvez, R., Martel, L., & Leroueil, S. (2010). Treatment of salted road runoffs using Typha latifolia, Spergularia canadensis, and Atriplex patula: a comparison of their salt removal potential. In Contaminated Sediments: Sustainable Management and Remediation. ASTM International. • Rozema, E. R., Gordon, R. J., & Zheng, Y. (2014). Plant species for the removal of Na+ and
- Cl-from Greenhouse Nutrient Solution. HortScience, 49(8), 1071-1075. • Artelia Eau et Environnement. 2018. MARCHE SUBSEQUENT N° 2018-SDAP-JL01 AU LOT 2 DE L'ACCORD-CADRE N°16P00165 Caractériser les possibilités d'évacuation d'un dépôt terrestre de sédiments de dragage
 - Région Occitanie, Parlement de la mer. 2015. Schéma régional des dragages en occitanie. Phase 1.





