



**HAL**  
open science

## **Arsenic and other trace elements in groundwaters and surface waters in the gold mining region of the Nigerien Liptako (Southwestern Niger)**

Oumar El Farouk Maman Illatou, Corinne Casiot, Marc Vinches, Eléonore Resongles, R. Freydier, Mylène Marie, Sophie Delpoux, Angelique Desoeuvre, Moussa Konaté, Boureima Ousmane

### ► To cite this version:

Oumar El Farouk Maman Illatou, Corinne Casiot, Marc Vinches, Eléonore Resongles, R. Freydier, et al.. Arsenic and other trace elements in groundwaters and surface waters in the gold mining region of the Nigerien Liptako (Southwestern Niger). *Environmental Earth Sciences*, 2022, 81 (23), pp.524. 10.1007/s12665-022-10639-8 . hal-03857742

**HAL Id: hal-03857742**

**<https://imt-mines-ales.hal.science/hal-03857742>**

Submitted on 29 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Arsenic and other trace elements in groundwaters and surface waters in the gold mining region of the Nigerien Liptako (Southwestern Niger)

Maman Illatou Oumar El Farouk<sup>1,2</sup> · Corinne Casiot<sup>1</sup> · Marc Vinches<sup>1</sup> · Eléonore Resongles<sup>1</sup> · Rémi Freydier<sup>1</sup> · Mylène Marie<sup>1</sup> · Sophie Delpoux<sup>1</sup> · Angélique Desoeuvre<sup>1</sup> · Moussa Konaté<sup>2</sup> · Boureima Ousmane<sup>2</sup>

## Abstract

The Nigerien Liptako in southwest Niger has gold mineralisations in Birimian formations (Paleoproterozoic) that have been mined for several decades. The objective of this study was to document the concentrations of arsenic and other toxic trace elements in groundwaters, river waters, suspended particulate matter and sediments and discuss the potential influence of artisanal gold mining on arsenic and trace elements enrichment. For this, forty-three groundwater samples (boreholes and wells) were analysed in mined and unmined areas. Surface waters, suspended particulate matters and sediments were sampled in three rivers at stations upstream and downstream from mining sites. Rock samples extracted from mine galleries were collected for mineralogical and geochemical characterization. In groundwater, arsenic concentrations ranged from 0.018 to 202 µg/L, exceeding drinking water guidelines (10 µg/L) in three artisanal gold mining areas. Manganese exhibited concentrations exceeding the potability limit in 23% of the samples, mainly in the Continental Terminal aquifer. In rivers, arsenic was the element most enriched in suspended particulate matter and sediments, but its concentration in water did not exceed 3 µg/L. Mineralogical analysis of rocks showed traces of arsenic in association with iron oxides, suggesting a release of arsenic in groundwater by desorption. The consumption of these waters rich in arsenic or manganese presents serious risks for health. These data represent the first large-scale investigation of arsenic and trace elements concentrations in groundwaters and surface waters in the Nigerien Liptako, where no or very scarce water quality measurements have been published yet.

**Keywords** Nigerien Liptako · Arsenic · Artisanal gold mining · Trace elements · Environment health

## Introduction

Gold mining is a potential source of high groundwater As concentration (Smedley and Kinniburgh 2002). The local disruption of chemical equilibrium in aquifer material generated by mine workings can lead to displacement of elements naturally present in the rocks, such as arsenic and trace metals. These elements are mobilized towards the water where they can reach concentrations harmful to the health of

consumers. In West Africa, gold mining has undergone significant development since the 1980s, with an acceleration in the last five years. The Liptako–Gourma (L.G.) region in West Africa straddles the borders shared by three countries, Niger, Mali and Burkina Faso. The area is rich in industrial and artisanal gold mines. Few studies investigated As concentration values in groundwater and surface water in gold mining areas from LG. In northern Burkina Faso, the presence of arsenic levels up to 1630 µg/L was shown in wells and boreholes in the Birimian rock aquifers in the locality of Ouahigouya. About 560,000 people are potentially exposed to groundwater arsenic contamination in this area (Smedley et al. 2007; Bretzler et al. 2017). In Mali, groundwater arsenic concentrations ranged from 0.8 to 132 µg/L in the village of Sanso, next to the Au Sanso mine, and it was accompanied by manganese at concentrations of 2.52 to 5200 µg/L (Bokar et al. 2020). These data suggest that As and trace

---

✉ Corinne Casiot  
corinne.casiot-marouani@umontpellier.fr

<sup>1</sup> HydroSciences Montpellier, Univ Montpellier, IRD, CNRS, IMT Mines Ales, Montpellier, France

<sup>2</sup> Département de Géologie, Faculté Des Sciences et Techniques, Université Abdou Moumouni, BP 10662, Niamey, Niger

elements enrichment in groundwaters in gold mining areas in the Liptako–Gourma region is of concern.

In the Nigerien Liptako, there has been no large-scale investigation to document arsenic and trace elements concentrations in water resources. Tankari et al. (2014) revealed the existence of significant water contamination by arsenic and trace elements in the vicinity of the Komabangou artisanal gold mining zone (Tillabéri, Niger). Wastewater from cyanide treatment exhibited extremely high concentrations of As, Cu, Mn, Ni, Pb and Zn (Tankari et al. 2014, 2019). In groundwater, average concentration values were 19 µg/L for As and 98 µg/L for Mn (Tankari et al. 2014). Apart from the Komabangou artisanal gold mining site, no survey of arsenic and other trace elements in the water resource has been carried out. The objective of the present study was to make a preliminary diagnosis of the levels of arsenic and trace metals in groundwaters, surface waters, river sediments and suspended particulate matter in gold mining areas in the Nigerien Liptako and to identify the source of these elements in water and the mobilization processes. For this, two sampling surveys were carried out in 2019 and arsenic and trace elements were analysed in water and solid samples. The two redox species of arsenic were also determined for a better insight into As mobilization processes. The source rocks were analysed by Scanning Electron Microscopy coupled to Energy Dispersive Spectroscopy (SEM–EDS) to find out the main mineral phases bearing arsenic and trace elements.

## Materials and methods

### Study area

#### General description

The study area is located in the southwestern part of Niger, next to the border with Burkina Faso and Mali, and lies between longitudes 0°30' W and 3°48' E, and latitudes 12°12' S and 15° N. It occupies the eastern part of the Nigerien Liptako and the western edge of the Iullemenden basin and covers the Nigerien part of the Niger River watershed with its tributaries on the right bank (Dargol, Sirba and Goroubi) and on the left bank (Kori of Ouallam, Dallol Bosso and Dallol Maouri) (Fig. 1a, b, c). The Liptako–Gourma (L.G.) zone covers an area of 370,000 km<sup>2</sup>. The Nigerien Liptako corresponds to the NE portion of the Man Dorsal (West African Craton). It mainly consists of the Birimian (Paleoproterozoic) formations. Gold has been mined at an artisanal level for decades in the area and there are currently over 100 artisanal gold mines in operation employing approximately 500,000 gold miners. Artisanal gold mining began in 1983, after a terrible drought and has increased in recent years with the arrival of refugees due to

insecurity in the Sahel. The extraction of gold in this area is considered to be a source of major environmental and health impacts (Barro-Traoré et al. 2008; Ochieng et al. 2010; Adama et al. 2012; Ouédraogo and Amyot 2013; Adama et al. 2017; Bretzler et al. 2017; Tankari 2014, 2019; Ibrahim et al. 2019, Bintou et al. 2020, Bokar et al. 2020), but its production contributes to the increase of the gross domestic product (GDP) and employment for many West African countries.

In the study area, the problem of water resource depletion has been increasing in recent years in relation with irregular rainfall, recurrent droughts in the Sahel, combined with an increasing demand for water, in relation with the rapid growth of the population (Ousmane et al. 2006). Degradation of water quality due to gold mining would further increase pressure on water resources.


### Geology

The geology of the study area consists of the Precambrian basement of the Liptako–Gourma to the west and the Iullemenden sedimentary basin to the east. The Nigerien Liptako consists of alternating greenstone belts and granitoid plutons oriented in a general NE–SW direction (Fig. 2a, NNW–SSE geological section of the study area (A–B)). The belt formations consist of metasediments, metavolcanosedimentary and metavolcanoplutonic formations (Machens 1973; Pouclet et al. 1990; Ama Salah 1996; Abdou et al. 1998; Soumaila 2000; Soumaila et al. 2004) which are composed of sandstone rocks (graywackes), mica-schists, shales, quartzschists, rhyolitic tuffs and breccias and green rocks (pyroxenites, amphibolites, epidotites, chlorite-schists, metabasalts, metagabbros). A variety of granitoid rocks including diorite, granodiorite and granites were intruded the greenstone belts, following the main structural trends (Dupuis et al. 1991; Cheilletz et al. 1994; Pons et al. 1995; Soumaila et al. 2005). The cover formations are formed of 5 to 50 m thick alterite, alluvium and colluvium.

The Continental terminal consists of the "Ader Douchi siderolithic series" (Ct<sup>1</sup>), the "lignite clay-sand series" (Ct<sup>2</sup>) and the "Middle Niger claystone series" (Ct<sup>3</sup>) (Greigert 1966). A geological section from Favreau (2000) is available in Fig. S1.

### Hydrogeology


The study area is characterized by discontinuous aquifers of the Liptako–Gourma basement (Fig. 2b, NNW–SSE hydrogeological section of the study area (C–D)), generalized aquifers of the Continental terminal formations (Fig. S1) and alluvial aquifers. The discontinuous aquifers are those of the alterites, the volcano-sedimentary series and the granite-gneiss formations (Ousmane 1988). The thickness



**Fig. 1** Location of sampling sites, artisanal gold mining sites, sampling grid, geology of the area and geological and hydrogeological cross sections (a) and zoom on the Dargol and Sirba watersheds showing river sampling sites (b) or rock and groundwater sampling sites (c)

of alteration varies from 10 to 20 m in the granite-gneissic domain (Guiraud and Travi 1990), and from 15 to 40 m in the shale and greenstone domain with a predominance of clay (Kamagaté et al. 2007). The recharge mechanism and groundwater origin of the fissured and fractured Liptako basement aquifers have been described through piezometric

methods and hydrogeochemistry (Babaye et al. 2018). The Continental terminal aquifers (Fig. S1) form three horizons recognized from bottom to top as Ct<sup>1</sup>, Ct<sup>2</sup>, and Ct<sup>3</sup> (Leduc and Karbo 1996). It is characterized by a weakness of the reserves in contact with the base and thickening of the aquifer series from the border of the Nigerien Liptako



**Fig. 2** NNW–SSE geological section of the Nigerien Liptako showing the succession of granitoid plutons and greenstone belts (a) and NNW–SSE hydrogeological section showing the alteration aquifer in the Ayorou pluton (b)

basement to the east of the study area. Isotopic analysis of the Ct aquifers has confirmed their current renewal (Leduc and Taupin 1997; Le Gal La Salle et al. 2001; Favreau et al. 2009). Their recharge is ensured by the direct infiltration of rainwater in the permeable outcrop areas and especially the emptying of water bodies (ponds) in the low-lying areas of the valleys, particularly in the Dallol Bosso and Kori of Ouallam sectors (Favreau 2000; Guéro 2003). Their annual recharge estimated through geochemical, hydrogeophysical and hydrodynamic approaches is 25 mm/year over the decades 1990–2000 (Favreau et al. 2009). The alluvial aquifers are contained in formations resulting from alluvial deposits of the Niger River or from tributary valleys.

## Sampling

### Sampling strategy

The sampling area covers the hydrographic basin of the Niger River in the Nigerien part, distributed on both sides of the two banks constituting the Niger River and its tributaries over an approximate area of 400 km × 200 km (80,000 km<sup>2</sup>). A preliminary study of the area according to its size, the types of anthropic activities that may release metallic pollutants into the environment, as well as the heterogeneity

of the geological formations oriented perpendicular to the direction of the Niger River was carried out. This study allowed the placement of a systematic sampling grid of 50 km × 50 km, superimposing the direction of the Niger River and perpendicular to the Sirba River in the N45 direction (Fig. 1). The aquifers involved are the discontinuous Liptako-Gourma basement aquifer (fractured or weathered) and alluvial aquifers on the right bank, and the Continental terminal (CT) aquifer on the left bank. On the right bank, the groundwater sampling points were chosen on the basis of the spatial distribution of artisanal gold mining sites, according to their proximity to the tributaries of the Niger River and considering their accessibility. On the left bank, proximity to the tributaries of the Niger River, notably the Kori of Ouallam and the Dalol Bosso, was mainly taken into consideration, as well as the size of the surrounding population.

The selection of surface water sampling points was guided by the morphology of the watershed and the spatial distribution of the gold-panning sites relative to the rivers. These rivers are the Dargol, the Sirba and the Goroubi. The northernmost Goroual River, which has been affected by intense gold-panning activities in the past, could not be sampled during the study period because of insecurity problems. The Dargol River lies south of a former large gold-panning site in Niger (Komabagou) and receives water from

surrounding sites. On the river Sirba, several placer gold mining sites have been identified. The Goroubi River is not directly involved in the artisanal exploitation of gold but is located downstream several former gold-panning sites such as Tangounga, on the border between Burkina and Niger. Four stations were selected on the Dargol river (Darg 1 to Darg 4), seven stations on the Sirba river (Sirb 1 to Sirb 7), and two stations on the Goroubi river (Goro 1 and Goro 2).

### Rock sampling

Samples of Nigerien Liptako rocks (basalt, granite, diorite, schist, quartz vein, quartzite and gneiss) were collected near major artisanal gold mining sites and at mineralized areas to examine the relationship between their mineralogical composition and trace metal concentrations in water (R1 to R8, Fig. 1).

### Water sampling

**Groundwater** Forty-three (43) water points were sampled for groundwater. Thirty samples were collected during the high-water periods between June and August 2019, and thirteen during the low water period in April and May in 2019. Depth of boreholes and wells ranged from 30 to 98 m on the right bank and down to 150 m on the left bank. The sampling was carried out on wells and boreholes regularly used for drinking water by the population. The water was collected in 1-L polyethylene bottles. The main physico-chemical parameters (pH, conductivity) were measured in situ using a calibrated multiparameter HACH®. Water was filtrated through 0.22 µm cellulose acetate disposable filter with a syringe, and distributed into polyethylene bottles for the determination of trace element (60 ml), major anions and cations (10 ml) and arsenic speciation (10 ml). The vials for trace element analysis and arsenic speciation were previously cleaned with 10% nitric acid (v:v) for 48 h and then rinsed three times with ultrapure water (Milli-Q®) in a clean room at the HydroSciences Montpellier laboratory. For alkalinity, vials (50 ml) were filled with unfiltered water to prevent air entry. The samples for trace element analysis were acidified with 60 µl of nitric acid (15 M HNO<sub>3</sub>, Suprapur, Merck). Samples for arsenic speciation were preserved with 100 µL of 8.7 M acetic acid solution and 100 µL of 50 g/L EDTA solution (Samanta and Clifford 2005). All water samples were kept at 4 °C during transport (cool box) and storage until analysis.

**River water** At each station on the rivers Sirba, Dargol and Goroubi, the main physico-chemical parameters (pH, conductivity) were measured and both water and surface sediments were sampled. Water (1 l) was collected in a polyethylene bottle and filtrated (250 ml to 500 ml) using a 500 ml

Nalgene filtration unit and manual vacuum pump on PVDF 0.22 µm filters. The filters were previously cleaned in 1% nitric acid, rinsed with ultrapure water, dried, weighed and preserved in a filter-box until use. Once filtrated, water was distributed into vials for trace elements, major anions and cations, arsenic speciation as described for groundwaters. For alkalinity, vials were filled with unfiltered water. The filter with suspended particulate matter (SPM) was recovered and stored in the filter-box, then air-dried in a desiccator and weighed before acid digestion.

### Sediment sampling

Fine surface sediments were collected after the rainy season (July 2019), while the waters begin to recede. They were sampled at each station in the river bed, under water in the main stream, using a spatula and stored in polypropylene vials, then frozen and freeze-dried before further processing.

## Sample processing and analysis

### Digestion procedure

Rocks and sediments were crushed in an agate mortar before acid digestion.

Samples were processed in a clean room class 10,000 at the HydroSciences laboratory. All consumable (vials, pipette tips...) was acid-cleaned before use and reagents were of Suprapur quality (Merck). Filter with SPM, sediments (50 mg) and rocks (50 mg) were digested in PTFE vials using a microwave oven (Ultrawave, Milestone) and a mixture of 15 M HNO<sub>3</sub> (3 ml), 9.5 M HCl (1 ml) and 22.6 M HF (1 ml). The temperature program was as follows: 5 min at 100 °C, then 5 min at 150 °C, and finally 10 min at 200 °C. The solution was evaporated to dryness at 70–80 °C. The residue was redissolved in 3 ml of 15 M HNO<sub>3</sub>, ultrasonicated for 20 min and heated at 100 °C on a hot plate during 10 min. After cooling, the solution was transferred in a 30 ml vial and completed with 27 mL of ultrapure water. This solution was diluted ten times with ultrapure water for ICP-MS analysis. To control the quality of analytical results, a certified reference material (NIST 2711a) and a blank of procedure were processed as for samples.

### Chemical and mineralogical analyses

Trace element concentrations in water, SPM, sediment and rock samples were determined by Inductively Coupled Plasma Mass Spectrometry (ICP-MS) on the AETE-ISO platform of OSU OREME (University of Montpellier) using routine procedures. Depending on the conductivity values, the water samples were diluted one, two or three times prior to analysis. A certified reference water

(SLRS-6) was analysed to check the quality of the analysis. Measured concentrations were within the uncertainty of the certified values for all studied elements (Al, As, B, Ba, Ca, Cd, Co, Cu, Cr, Fe, K, Mg, Mn, Na, Ni, Pb, Zn) and the analytical error (relative standard deviation) was better than 5%, except for Cd (8%). In addition, accuracy was within 10% of the certified values for the reference material NIST 2711a for all elements (except 19% for B) and blank contribution was negligible (<1% for all elements, except for Ba, Cr, Pb <5%) confirming that no contamination occurred during the digestion procedure.

Redox arsenic species were analysed by anion-exchange chromatography (25 cm, 4.1 mm inner diameter Hamilton PRP-X100 column) using a Spectra Device SCM1000 solvent delivery pumps coupled to ICP-MS (Héry et al. 2014). Major anions and cations in water samples were analysed by DIONEX ICS-1000 ion chromatography at HydroSciences Montpellier laboratory. Alkalinity was determined by acid titration. The analysed compounds are  $\text{HCO}_3^-$ ,  $\text{F}^-$ ,  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{PO}_4^{3-}$  for the anions and  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$  and  $\text{NH}_4^+$  for the cations. The reliability of the results of the major element analyses was validated after the calculation of the ionic balance values ( $-7 < \text{BA} < +7$ ). The calculated values are considered acceptable (<10%). The software used for the data analysis (Piper diagram), is DIAGRAMME from the University of Avignon.

Mineralogical analyses were performed by scanning electron microscopy coupled with Energy Dispersive Spectroscopy (SEM-EDS) at the C2MA laboratory of IMT Mines Alès, after the preparation of polished thin slides.

## Enrichment factors

Trace element enrichment factors (EF) in suspended particulate matter and sediments were calculated according to the following equation:

$$F = \frac{\left(\frac{[M]}{[Al]}\right)_{\text{sample}}}{\left(\frac{[M]}{[Al]}\right)_{\text{av.crust}}},$$

where  $\left(\frac{[M]}{[Al]}\right)_{\text{sample}}$  represents the ratio between element concentration and aluminium concentration in the sample and  $\left(\frac{[M]}{[Al]}\right)_{\text{av.crust}}$  represents the ratio between average element concentration and aluminium concentration in the earth surface continental crust (average Upper Continental Crust, Taylor and McLennan 1995). Due to the lack of published data for the geochemical composition of sediments in the study region, the upper continental crust value is used as reference in the calculation of the enrichment factor even though it may not reflect the regional background values.

## Statistical treatment

The statistical analyses were performed with the R free software. Groundwater trace elements concentrations data were analysed by non-parametric test Mann–Whitney; the  $p$  values were obtained at the level of  $p < 0.05$ .

## Results

### Rocks

Table 1 shows the concentrations of arsenic and other trace elements in the different types of rocks collected in the gold exploitation zones and the main mineralogy inferred from SEM-EDS analysis (Fig. S2). The selected rocks are associated with gold mineralisations and hence represent the material extracted and processed on gold-panning sites that represents a potential source of geogenic metallic contaminants for the surrounding hydrosystems.

The As content of the rocks ranged between 0.4 and 8.3  $\mu\text{g/g}$  (Table 1). These concentrations were in the same order of magnitude than the average values for basalt (2.3  $\mu\text{g/g}$ ), granite (1.3  $\mu\text{g/g}$ ), diorite (1  $\mu\text{g/g}$ ), schist (1.1  $\mu\text{g/g}$ ), quartzite (5.5  $\mu\text{g/g}$ ) and gneiss (1.1  $\mu\text{g/g}$ ) reported by Smedley and Kinniburgh (2002). Arsenopyrite (FeAsS) was not detected by SEM in the rock samples of the studied area. Arsenic was present as traces (~0.8%) in Fe/Mn oxides in the schist sample from Doumba (Fig. S2).

The most remarkable result was the high Mn (85 mg/g) and Ba (27 mg/g) contents in the sample of the quartz vein (quartz diorite) from the Torodi pluton (Goroubi watershed) (Table 1). This was consistent with the detection of Mn oxide and Ba carbonate by SEM (Fig. S2). Barium was also detected as barite ( $\text{BaSO}_4$ ) in the basalt sample from Sirba A (village of Talé), despite the low Ba content (84  $\mu\text{g/g}$ ) in this rock (Table 3). The concentrations of Co, Mn, Ni, and Cr were enriched relatively to the average composition of the upper continental crust for most types of rocks from the studied area while Ba, Cd, Cu concentrations exceeded average values in some samples. There was no Co, Ni, or Cr minerals detected by SEM. These elements were detected as traces (~2% Co; ~3% Ni, ~0.4% Cr) in association with Fe/Mn oxides in the schist sample from Doumba (Fig. S2). Several metal sulphides were detected by SEM in the magmatic rock samples of the study area: pyrite ( $\text{FeS}_2$ ), chalcopyrite ( $\text{CuFeS}_2$ ) and other sulphide minerals consisting of pyrrotite ( $\text{Fe}_{1-x}\text{S}$ ), chalcocite ( $\text{Cu}_2\text{S}$ ) and pentlandite ( $(\text{Fe}, \text{Ni})\text{S}$ ) were detected in the basalt sample (Fig. S2).

**Table 1** Concentration (in % or µg/g) of major and trace elements and mineral phases in selected rock samples

| Sample name  | Watershed | Type of rock                 | Al (%)    | Ca (%)    | Fe (%)    | K (%)                        | Na (%)                 | Mg (%)                   | As (µg/g)                               | B (µg/g) | Ba (µg/g) | Cd (µg/g) | Co (µg/g) | Cr (µg/g) | Cu (µg/g) | Mn (µg/g) |
|--|-----------|------------------------------|-----------|-----------|-----------|------------------------------|------------------------|--------------------------|---|----------|-----------|-----------|-----------|-----------|-----------|-----------|
| R1   | Sirba     | Basalt                       | 8.35      | 9.05      | 6.87      | 0.123                        | 2.03                   | 4.82                     | 0.368                                   | 8.66     | 82.6      | 0.034     | 67.5      | 209       | 3.27      | 833       |
| R2   | Sirba     | Granite                      | 12.4      | 5.79      | 5.34      | 3.12                         | 5.63                   | 1.71                     | 1.08                                    | 4.64     | 1232      | 0.066     | 121       | 61.3      | 12.1      | 928       |
| R3   | Dargol    | Schist                       | 4.72      | 0.050     | 7.26      | 0.008                        | 0.006                  | 11.5                     | 8.28                                    | 1.01     | 15.2      | 0.014     | 120       | 3425      | 20.7      | 758       |
| R4   | Dargol    | Quartzite                    | 0.055     | 0.034     | 0.071     | 0.007                        | 0.008                  | 0.114                    | 0.423                                   | 2.36     | 3.32      | 0.012     | 195       | 26.2      | 1.32      | 11.6      |
| R5   | Sirba     | Basalt                       | 7.47      | 8.70      | 7.47      | 0.058                        | 1.16                   | 4.34                     | 1.02                                    | 5.36     | 83.7      | 0.098     | 85.5      | 187       | 86.1      | 1449      |
| R6   | Sirba     | Diorite                      | 9.24      | 6.85      | 6.49      | 1.14                         | 1.92                   | 4.40                     | 2.56                                    | 11.2     | 441       | 0.125     | 70.8      | 313       | 64.5      | 1145      |
| R7   | Goroubi   | Quartz diorite (quartz vein) | 0.647     | 0.099     | 0.447     | 0.043                        | 0.027                  | 0.024                    | 2.24                                    | 1.09     | 26,898    | 0.664     | 740       | 22.6      | 143       | 84,768    |
| R8   | Dargol    | Gneiss                       | 1.66      | 10.0      | 3.72      | 0.128                        | 0.250                  | 5.41                     | 1.49                                    | 1.74     | 75.0      | 0.591     | 72.4      | 1297      | 1.6       | 1268      |
| Average composition of the upper continental crust** |           |                              | 8.04      | 3.00      | 3.50      | 2.80                         | 2.89                   | 1.33                     | 1.5                                     | 15       | 550       | 0.098     | 10        | 35        | 25        | 600       |
| Sample name  | Watershed | Type of rock                 | Ni (µg/g) | Pb (µg/g) | Zn (µg/g) | Published As concentrations* | Sulphides              | Mineralogy oxides        | Other                                   |          |           |           |           |           |           |           |
| R1   | Sirba     | Basalt                       | 100       | 0.512     | 29.4      | 2.3                          | Not analysed           |                          |   |          |           |           |           |           |           |           |
| R2   | Sirba     | Granite                      | 37.7      | 9.39      | 81.9      | 1.3                          | Not analysed           |                          |   |          |           |           |           |           |           |           |
| R3   | Dargol    | Schist                       | 1523      | 0.648     | 65.4      | 1.1                          |                        | Co-, Ti-, Fe-, Mn-oxides | Quartz                                  |          |           |           |           |           |           |           |
| R4   | Dargol    | Quartzite                    | 12.5      | 0.220     | 2.36      | 5.5                          | Not analysed           |                          |   |          |           |           |           |           |           |           |
| R5   | Sirba     | Basalt                       | 113       | 0.631     | 67.6      | 2.3                          | Cu-, Ni-, Fe- sulphide |                          | Quartz, Ba sulphate, Ti-Aluminosilicate |          |           |           |           |           |           |           |
| R6   | Sirba     | Diorite                      | 80.9      | 5.27      | 71.9      | 1                            | Not analysed           |                          |   |          |           |           |           |           |           |           |
| R7   | Goroubi   | Quartz diorite (quartz vein) | 99.6      | 5.25      | 78.7      | 5.5/1                        |                        | Ce-, Mn-oxide            | Ba carbonate                            |          |           |           |           |           |           |           |
| R8   | Dargol    | Gneiss                       | 613       | 3.34      | 81.3      | 1                            | Not analysed           |                          |   |          |           |           |           |           |           |           |
| Average composition of the upper continental crust** |           |                              | 20        | 20        | 71        |                              |                        |                          |   |          |           |           |           |           |           |           |

\*Smedley et al. 2002

\*\*Taylor and McLennan (1995)



## Groundwater

The main physico-chemical parameters and concentrations of major and trace elements in groundwater are presented in Table 2.

Most groundwater samples showed calcic and magnesian bicarbonate facies, for the two geological environments (Nigerien Liptako and CT<sup>3</sup>) (Table 2, Fig. S3). A few samples in the CT<sup>3</sup> geologic environment exhibited a sodium potassium chloride or a calcium magnesium chloride and sulphate composition type (Table 2). Ten samples had nitrate concentration above drinking water guidelines (Nigerien standard: 45 mg/L), eight in LG geological environment, two in CT geological environment. The origin of nitrate was attributed in the study area to the use of fertilizers and manure for agriculture and to human settlements (Salihou-Djari et al. 2018).

Trace element concentrations in groundwaters were highly variable within each geological environment (Fig. 3). Arsenic concentrations exceeded drinking water guidelines (10 µg/L) in four samples, all in LG geological environment (Fig. 4, Table 2). The two samples with the highest As concentrations corresponded to wells in Nbang (203 µg/L) and Boulounjouna (190 µg/L) on the Sirba watershed. Both were located in a gold mining area and have been dug decades ago for gold extraction; the water recovered from these wells during dewatering is used for drinking water purpose by the surrounding population. Comparatively, the well in Sefa Moussa exploited since 2017 for gold extraction exhibited lower As concentration (2.5 µg/L). The two samples with 16 µg/L of As corresponded to boreholes close (< 1 km) to the old (> 40 years) Komabangou mining site; they have been drilled specifically for drinking water supply. In CT, only one sample exceeded 1 µg/L (borehole from Sona, 2.1 µg/L of As). Considering all data, arsenic concentrations were significantly higher ( $p < 0.05$ ) in LG geological environment than in CT geological environment. The oxidized arsenate species predominated in all groundwater samples analysed, except in the well from Sefa Moussa, where the reduced species arsenite (As(III)) predominated at low concentration (2.5 µg/L).

Manganese concentrations exceeded guidelines in ten samples, both in LG and CT geological environments (Fig. 5, Table 2). However, considering all data, Mn concentrations were higher ( $p < 0.05$ ) in CT geological environment than in LG geological environment (Fig. 3). Barium concentrations exceeded Nigerien drinking water guidelines (100 µg/L) in nineteen samples, both in LG and CT geological environments (Table 2); however, only one sample exceeded WHO guidelines (1300 µg/L). Two samples exceeded boron drinking water guidelines and one sample exceeded nickel guidelines, all in CT geological environment (Table 2). Other trace elements Al, Fe, Co, Cu, Zn, Cd

and Pb did not exceed drinking water guidelines (Table 2). No strong correlation was found, neither between trace element concentrations and hydrochemical parameters, nor among trace elements.

## Rivers

### Water

Surface water of the Rivers Sirba, Dargol and Goroubi exhibited calcic and magnesian bicarbonate hydrochemical facies (Fig. S3). One sample on the river Sirba (village Larba) contained more chloride and sulphate (Table 2).

The concentrations of arsenic and other trace elements in the dissolved phase were compared with published values for the Niger River and to world rivers averages (Table 2). Arsenic concentrations were at maximum five times higher than world rivers average (no data available for the river Niger); they were higher in the River Sirba ( $[As]_{\text{average}} = 1.5 \pm 0.9 \mu\text{g/L}$ ,  $n = 7$ ) than in the Rivers Dargol ( $[As]_{\text{average}} = 0.3 \pm 0.2 \mu\text{g/L}$ ,  $n = 4$ ) and Goroubi ( $[As] = 0.13\text{--}0.19 \mu\text{g/L}$ ,  $n = 2$ ). In the river Sirba, arsenic concentration increased between stations Sirb6 and Sirb3, located upstream and downstream from a gold-panning area. Arsenite (As(III)) represented more than 70% of total As concentration in the river Sirba. In the River Dargol, arsenate predominated at low concentration (0.58 µg/L at station Darg 1).

Manganese, Co, Zn and Ba were the elements most enriched in the dissolved phase compared to the River Niger or to world river averages, up to a factor 323 for Mn, 50 for Co, 26 for Zn and 29 for Ba.

### Suspended particulate matter and sediments

Concentrations of arsenic and other trace elements in suspended particulate matter and sediments are presented in Table 3, together with enrichment factors relatively to upper continental crust. Concentrations in SPM were generally similar or lower than world river averages, except Cd at the most downstream station (Sirb 1) in the river Sirba (Table 3). They were highly correlated ( $r^2 > 0.7$ ) to the content of Al (for Co, Cr, Cu, Fe, Ni, Pb, Zn) and Fe (for B, Ba, Co, Cr, Cu, Ni, Pb, Zn). The correlation ( $r^2 > 0.7$ ) was also found in sediments between some of the trace elements and Al (for Ni, Pb, Zn) or Fe (for As, Co, Cr, Cu).

Arsenic exhibited the highest enrichment factors in both SPM and sediments, with maximum values in the river Sirba (EF = 6 to 14 in SPM and EF = 10 to 59 in sediments). The highest EF (EF = 59) was reached for sediments at station Sirb3, located downstream from a gold-panning area. For other trace elements, enrichment factors were generally lower than 4, except locally for Cd (Sirb 1) and Mn (Sirb 2)

**Table 2** Main physico-chemical parameters and concentrations of major and trace elements in groundwater (wells and boreholes) and surface water. For arsenic, percentage of arsenite and arsenate are mentioned, when available

| Sample name | Latitude | Longitude | Sampling date (wet (W)/dry (D) period) | Watershed       | Geological environment | Structure | Water use       | Aquifer type        | Geological formation       | Depth (m) |
|-------------|----------|-----------|--|-----------------|------------------------|-----------|-----------------|---------------------|----------------------------|-----------|
| Groundwater |          |           |  |                 |                        |           |                 |                     |                            |           |
| P1          | 13.61857 | 1.57649   | 17/04/2019, D                          | Sirba           | LG                     | Well      | Gold extraction | Alterites           | Green rocks                | 34        |
| P2          | 14.08278 | 1.06320   | 13/06/2019, D                          | Dargol          | LG                     | Well      | Drinking water  | Alterites           | Altered schists            | 60        |
| P3          | 13.67982 | 1.61183   | 13/06/2019, D                          | Sirba           | LG                     | Well      | Gold extraction | Alterites           | Green rocks                | 40        |
| P4          | 13.45657 | 1.17384   | 17/07/2019, W                          | Sirba           | LG                     | Well      | Gold extraction | Alterites           | Green rocks                | 50        |
| P5          | 13.42506 | 1.17241   | 17/07/2019, W                          | Sirba           | LG                     | Well      | Drinking water  | Alterites           | Green rocks                | 10        |
| P6          | 13.40817 | 1.22886   | 17/07/2019, W                          | Sirba           | LG                     | Well      | Drinking water  | Alterites           | Green rocks                | 20        |
| P7          | 14.11846 | 1.15282   | 22/09/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Green rocks                | 67        |
| P8          | 14.00492 | 1.13031   | 22/09/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Fractures           | Fractured granite          | 67        |
| P9          | 13.73522 | 2.59041   | 09/05/2019, D                          | Dallol Bosso    | CT                     | Well      | Drinking water  | CT3                 | CT                         | 28        |
| P10         | 13.30080 | 3.35514   | 09/05/2019, D                          | Dallol Maouri   | CT                     | Borehole  | Drinking water  | CT3                 | CT                         | 35        |
| P11         | 14.95680 | 2.00260   | 23/08/2019, W                          | Kori of Ouallam | CT                     | Well      | Drinking water  | CT3                 | CT                         | 60        |
| P12         | 13.54170 | 2.24770   | 08/09/2019, W                          | Kori of Ouallam | CT                     | Well      | Drinking water  | CT3                 | CT                         | 39        |
| P13         | 13.68700 | 2.69720   | 08/09/2019, W                          | Dallol Bosso    | CT                     | Well      | Drinking water  | CT3                 | CT                         | 40        |
| P14         | 13.04690 | 2.54340   | 04/09/2019, W                          | Niger river     | CT                     | Borehole  | Drinking water  | CT3                 | CT                         | 50        |
| F1          | 13.80142 | 1.35662   | 07/05/2019, D                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Fractured granite          | 48        |
| F2          | 13.90552 | 0.80816   | 24/04/2019, D                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Fractured granite          | 72        |
| F3          | 12.75132 | 2.17564   | 08/05/2019, D                          | Tapoa river     | LG                     | Borehole  | Drinking water  | Alterites           | Fractured granite          | 46        |
| F4          | 13.37128 | 2.25026   | 09/05/2019, D                          | Niger river     | LG                     | Borehole  | Drinking water  | Alterites           | Fractured granite          | 88        |
| F5          | 14.00770 | 0.84406   | 25/04/2019, D                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Green rocks                | 60        |
| F6          | 14.08214 | 1.05564   | 13/06/2019, D                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Altered schists            | 70        |
| F7          | 13.58624 | 1.34927   | 17/07/2019, W                          | Sirba           | LG                     | Borehole  | Drinking water  | Alterites/Fractures | Fractured granite          | 64        |
| F8          | 12.84236 | 1.68933   | 17/07/2019, W                          | Goroubi         | LG                     | Borehole  | Drinking water  | Fractures           | Fractured granite          | 38        |
| F9          | 13.12687 | 1.79868   | 17/07/2019, W                          | Goroubi         | LG                     | Borehole  | Drinking water  | Fractures           | Basement                   | 59        |
| F10         | 14.43487 | 1.27655   | 07/08/2019, W                          | Niger river     | LG                     | Borehole  | Drinking water  | Fractures           | Fractured gneiss           | 57        |
| F11         | 14.69694 | 0.96663   | 07/08/2019, W                          | Niger river     | LG                     | Borehole  | Drinking water  | Fractures           | Fractured amphibolite      | 50        |
| F12         | 13.92861 | 1.02971   | 25/08/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Green rocks                | 90        |
| F13         | 14.19542 | 0.96625   | 25/08/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Fractures           | Fractured granite          | 55        |
| F14         | 14.19206 | 0.92973   | 25/08/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Fractures           | Fractured granite          | 57        |
| F15         | 14.05856 | 1.06229   | 16/07/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Green rocks                | 40        |
| F16         | 14.17848 | 1.03224   | 25/08/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Basement/Green rocks limit | 52        |
| F17         | 13.97690 | 1.55150   | 10/09/2019, W                          | Niger river     | CT                     | Borehole  | Drinking water  | CT3                 | CT                         | 75        |
| F18         | 14.15232 | 1.13172   | 22/09/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Alterites           | Green rocks                | 65        |
| F19         | 14.00009 | 1.18231   | 22/09/2019, W                          | Dargol          | LG                     | Borehole  | Drinking water  | Fractures           | Fractured granite          | 48        |

**Table 2** (continued)

| Sample name  | Latitude              | Longitude | Sampling date (wet (W)/dry (D) period) | Watershed               | Geological environment | Structure             | Water use              | Aquifer type            | Geological formation   | Depth (m)             |                                      |                                     |                                     |                                      |      |
|--------------|-----------------------|-----------|--|-------------------------|------------------------|-----------------------|------------------------|-------------------------|------------------------|-----------------------|--------------------------------------|-------------------------------------|-------------------------------------|--------------------------------------|------|
| F20          | 13.73399              | 2.58975   | 09/05/2019, D                          | Dallol Bosso            | CT                     | Borehole              | Drinking water         | CT2                     | CT                     | 154                   |                                      |                                     |                                     |                                      |      |
| F21          | 13.74667              | 2.10003   | 29/08/2019, W                          | Kori of Ouallam         | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 90                    |                                      |                                     |                                     |                                      |      |
| F22          | 14.67630              | 1.95355   | 23/08/2019, W                          | Kori of Ouallam         | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 60                    |                                      |                                     |                                     |                                      |      |
| F23          | 12.73883              | 2.86820   | 22/09/2019, W                          | Dallol Bosso            | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 30                    |                                      |                                     |                                     |                                      |      |
| F24          | 14.13664              | 2.13246   | 29/08/2019, W                          | Kori of Ouallam         | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 64                    |                                      |                                     |                                     |                                      |      |
| F25          | 12.36859              | 2.91603   | 30/08/2019, W                          | Niger river             | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 54                    |                                      |                                     |                                     |                                      |      |
| F26          | 14.45816              | 2.03066   | 23/08/2019, W                          | Kori of Ouallam         | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 49                    |                                      |                                     |                                     |                                      |      |
| F27          | 13.27214              | 2.95275   | 19/08/2019, W                          | Dallol Bosso            | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 30                    |                                      |                                     |                                     |                                      |      |
| F28          | 12.99932              | 3.25716   | 19/08/2019, W                          | Dallol Maouri           | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 98                    |                                      |                                     |                                     |                                      |      |
| F29          | 12.80400              | 3.63700   | 28/12/2019, D                          | Dallol Maouri           | CT                     | Borehole              | Drinking water         | CT3                     | CT                     | 57                    |                                      |                                     |                                     |                                      |      |
| Sample name  | Conduc-tivity (µS/cm) | T°C       | pH                                     | Ca <sup>2+</sup> (mg/L) | Cl <sup>-</sup> (mg/L) | K <sup>+</sup> (mg/L) | Na <sup>+</sup> (mg/L) | Mg <sup>2+</sup> (mg/L) | Br <sup>-</sup> (mg/L) | F <sup>-</sup> (mg/L) | HCO <sub>3</sub> <sup>-</sup> (mg/L) | NO <sub>3</sub> <sup>-</sup> (mg/L) | NO <sub>2</sub> <sup>-</sup> (mg/L) | SO <sub>4</sub> <sup>2-</sup> (mg/L) |      |
| Ground-water | P1                    | 783       | 32                                     | 7.54                    | 43.8                   | 7.01                  | 1.24                   | 43.7                    | 58.5                   | <0.02                 | 0.28                                 | 497                                 | 13.5                                | <0.08                                | 53.8 |
|              | P2                    | 473       | 32                                     | 6.80                    | 53.2                   | 6.76                  | 3.07                   | 20.8                    | 15.0                   | <0.02                 | 0.37                                 | 213                                 | 29.4                                | n.d                                  | 7.51 |
|              | P3                    | 828       | 24                                     | 7.10                    | 104                    | 7.91                  | 0.88                   | 29.3                    | 36.7                   | <0.02                 | 0.21                                 | 460                                 | 88.1                                | n.d                                  | 40.6 |
|              | P4                    | 1003      | 32                                     | 7.54                    | 12.5                   | 17.3                  | 5.75                   | 18.9                    | 54.4                   | <0.02                 | <0.08                                | n.d                                 | 1.11                                | <0.08                                | 19.2 |
|              | P5                    | 86        | 30                                     | 5.81                    | 1.74                   | 0.75                  | 0.26                   | 0.68                    | 0.38                   | <0.02                 | <0.08                                | 29.2                                | 0.84                                | <0.08                                | 0.47 |
|              | P6                    | 69        | 34                                     | 5.92                    | 6.50                   | 6.12                  | 8.31                   | 1.85                    | 2.40                   | <0.02                 | 0.14                                 | 28.4                                | 8.62                                | <0.08                                | 2.46 |
|              | P7                    | 117       | 20                                     | 5.81                    | 11.6                   | 3.20                  | 3.33                   | 3.50                    | 3.19                   | <0.02                 | <0.08                                | 31.7                                | 11.5                                | <0.08                                | 14.2 |
|              | P8                    | 597       | 19                                     | 6.47                    | 60.3                   | 3.65                  | 1.00                   | 42.0                    | 24.4                   | <0.02                 | <0.08                                | 291                                 | 39.8                                | <0.08                                | 71.2 |
|              | P9                    | 196       | 32                                     | 5.99                    | 21.2                   | 4.76                  | 1.58                   | 14.0                    | 1.78                   | <0.02                 | 0.05                                 | 62.4                                | 43.2                                | 0.21                                 | 1.70 |
|              | P10                   | 211       | 33                                     | 6.19                    | 5.25                   | 3.46                  | 7.38                   | 29.1                    | 5.89                   | <0.02                 | 0.24                                 | 110                                 | <0.04                               | <0.08                                | 16.6 |
|              | P11                   | 405       | 31                                     | 5.42                    | 30.9                   | 7.86                  | 3.13                   | 21.6                    | 11.7                   | 0.06                  | 0.16                                 | 12.1                                | 136                                 | 0.26                                 | 11.9 |
|              | P12                   | 292       | 31                                     | 6.27                    | 36.0                   | 12.8                  | 3.78                   | 10.6                    | 6.34                   | <0.02                 | 0.38                                 | 177                                 | 0.20                                | <0.08                                | 0.82 |
|              | P13                   | 198       | 31                                     | 5.96                    | 14.2                   | 7.01                  | 6.22                   | 18.6                    | 4.00                   | <0.02                 | <0.08                                | 63.6                                | 41.9                                | <0.08                                | 3.49 |
|              | P14                   | 58        | 31                                     | 5.60                    | 3.20                   | 1.17                  | 0.55                   | 8.84                    | 0.50                   | <0.02                 | <0.08                                | 12.4                                | 22.2                                | <0.08                                | 0.58 |
|              | F1                    | 576       | 32                                     | 6.51                    | 49.2                   | 18.8                  | 0.80                   | 69.8                    | 8.71                   | 0.15                  | 0.43                                 | 321                                 | 25.9                                | 0.31                                 | 12.6 |
|              | F2                    | 638       | 34                                     | 6.65                    | 45.0                   | 9.29                  | 3.43                   | 82.3                    | 15.9                   | 0.17                  | 0.65                                 | 393                                 | 13.8                                | 0.18                                 | 17.0 |
|              | F3                    | 522       | 33                                     | 6.89                    | 47.7                   | 3.00                  | 2.69                   | 32.2                    | 25.6                   | <0.02                 | 0.15                                 | 337                                 | 0.08                                | <0.08                                | 26.8 |
|              | F4                    | 604       | 36                                     | 6.50                    | 43.1                   | 23.9                  | 4.82                   | 45.6                    | 29.5                   | <0.02                 | 0.29                                 | 317                                 | 0.12                                | <0.08                                | 46.7 |
|              | F5                    | 935       | 34                                     | 6.46                    | 92.0                   | 16.5                  | 4.39                   | 65.7                    | 34.1                   | 0.29                  | 0.19                                 | 289                                 | 77.5                                | <0.08                                | 195  |
|              | F6                    | 742       | 36                                     | 7.43                    | 67.6                   | 10.6                  | 1.37                   | 69.6                    | 23.3                   | <0.02                 | 0.31                                 | 460                                 | 42.3                                | n.d                                  | 33.8 |

**Table 2** (continued)

| Sample name  | Conductivity (µS/cm) | T°C       | pH                            | Ca <sup>2+</sup> (mg/L)     | Cl <sup>-</sup> (mg/L) | K <sup>+</sup> (mg/L) | Na <sup>+</sup> (mg/L) | Mg <sup>2+</sup> (mg/L) | Br <sup>-</sup> (mg/L) | F <sup>-</sup> (mg/L) | HCO <sub>3</sub> <sup>-</sup> (mg/L) | NO <sub>3</sub> <sup>-</sup> (mg/L) | NO <sub>2</sub> <sup>-</sup> (mg/L) | SO <sub>4</sub> <sup>2-</sup> (mg/L) |           |       |
|--------------|----------------------|-----------|-------------------------------|-----------------------------|------------------------|-----------------------|------------------------|-------------------------|------------------------|-----------------------|--------------------------------------|-------------------------------------|-------------------------------------|--------------------------------------|-----------|-------|
| F7           | 469                  | 30        | 6.5                           | n.d                         | 44.0                   | 4.93                  | 27.3                   | 15.1                    | n.d                    | n.d                   | 82.3                                 | 93.0                                | n.d                                 | 4.22                                 |           |       |
| F8           | 487                  | 28        | 6.59                          | 53.0                        | 2.52                   | 3.21                  | 21.6                   | 19.4                    | <0.02                  | <0.08                 | 322                                  | 4.92                                | <0.08                               | 4.80                                 |           |       |
| F9           | 344                  | 28        | 8.68                          | 25.0                        | 7.86                   | 6.64                  | 15.6                   | 16.6                    | <0.02                  | 0.49                  | 95.0                                 | 31.0                                | <0.08                               | 2.73                                 |           |       |
| F10          | 329                  | 31        | 6.19                          | 36.1                        | 6.03                   | 2.37                  | 20.2                   | 9.15                    | <0.02                  | 0.15                  | 194                                  | 17.3                                | <0.08                               | 5.40                                 |           |       |
| F11          | 1095                 | 34        | 6.56                          | 98.9                        | 35.7                   | 1.92                  | 94.1                   | 35.1                    | <0.02                  | <0.08                 | 268                                  | 321                                 | <0.08                               | 21.4                                 |           |       |
| F12          | 498                  | 28        | 6.89                          | 46.9                        | 2.62                   | 3.86                  | 37.8                   | 20.7                    | <0.02                  | 0.54                  | 350                                  | 11.1                                | <0.08                               | 4.53                                 |           |       |
| F13          | 847                  | 28        | 7.38                          | 73.8                        | 18.7                   | 8.83                  | 24.2                   | 54.1                    | <0.02                  | <0.08                 | 419                                  | 111                                 | <0.08                               | 23.0                                 |           |       |
| F14          | 672                  | 28        | 6.96                          | 45.4                        | 6.30                   | 6.16                  | 76.1                   | 15.3                    | <0.02                  | <0.08                 | 465                                  | 1.00                                | <0.08                               | 11.5                                 |           |       |
| F15          | 448                  | 28        | 6.96                          | 44.3                        | 6.13                   | 3.40                  | 25.3                   | 16.1                    | <0.02                  | 0.48                  | 150                                  | 30.6                                | <0.08                               | 9.64                                 |           |       |
| F16          | 470                  | 28        | 6.96                          | 55.9                        | 5.61                   | 4.47                  | 14.4                   | 23.2                    | <0.02                  | <0.08                 | 333                                  | 6.89                                | <0.08                               | 6.14                                 |           |       |
| F17          | 963                  | 33        | 6.96                          | 41.7                        | 25.2                   | 1.55                  | 178                    | 15.9                    | <0.02                  | 1.49                  | 524                                  | 47.4                                | <0.08                               | 43.5                                 |           |       |
| F18          | 926                  | 21        | 6.59                          | 102                         | 26.6                   | 3.41                  | 69.5                   | 28.8                    | <0.02                  | <0.08                 | 404                                  | 159                                 | <0.08                               | 20.7                                 |           |       |
| F19          | 648                  | 20        | 6.32                          | 65.5                        | 16.2                   | 2.04                  | 28.2                   | 27.4                    | <0.02                  | <0.08                 | 187                                  | 144                                 | <0.08                               | 26.3                                 |           |       |
| F20          | 1337                 | 33        | 7.11                          | 54.3                        | 121                    | 10.3                  | 209                    | 25.5                    | 0.62                   | 1.77                  | 339                                  | 0.54                                | <0.08                               | 254                                  |           |       |
| F21          | 64                   | 31        | 5.35                          | 4.79                        | 2.86                   | 4.26                  | 2.48                   | 1.37                    | <0.02                  | <0.08                 | 4.64                                 | 14.1                                | <0.08                               | 8.76                                 |           |       |
| F22          | 524                  | 29        | 6.24                          | 50.3                        | 5.87                   | 3.71                  | 13.9                   | 13.7                    | <0.02                  | 0.29                  | 202                                  | 0.68                                | <0.08                               | 82.5                                 |           |       |
| F23          | 1142                 | 31        | 5.24                          | 15.3                        | 9.88                   | 1.03                  | 12.9                   | 4.49                    | <0.02                  | <0.08                 | 8.42                                 | 24.4                                | <0.08                               | 4.76                                 |           |       |
| F24          | 1275                 | 31        | 6.96                          | 55.8                        | 121                    | 9.53                  | 199                    | 26.4                    | <0.02                  | 1.01                  | 181                                  | 2.35                                | <0.08                               | 242                                  |           |       |
| F25          | 70                   | 31        | 5.79                          | 3.42                        | 0.86                   | 1.61                  | 1.08                   | 0.61                    | <0.02                  | <0.08                 | 22.3                                 | 3.55                                | <0.08                               | 0.71                                 |           |       |
| F26          | 817                  | 28        | 6.42                          | 80.5                        | 53.3                   | 7.56                  | 65.6                   | 24.7                    | 0.36                   | 0.41                  | 263                                  | 0.44                                | <0.08                               | 164                                  |           |       |
| F27          | 1605                 | 31        | 5.96                          | 62.3                        | 252                    | 71.2                  | 183                    | 19.3                    | <0.02                  | <0.08                 | 93.4                                 | 162                                 | <0.08                               | 141                                  |           |       |
| F28          | 72                   | 31        | 5.57                          | 3.10                        | 0.97                   | 0.54                  | 1.71                   | 0.78                    | <0.02                  | <0.08                 | 12.9                                 | 3.34                                | <0.08                               | 0.95                                 |           |       |
| F29          | 417                  | 25        | 7.60                          | 7.80                        | 23.9                   | 5.10                  | 79.6                   | 5.00                    | 0.17                   | 1.61                  | 223                                  | <0.04                               | <0.08                               | <0.50                                |           |       |
| Sample name  | Al (µg/L)            | As (µg/L) | As(III)/As <sub>tot</sub> (%) | As(V)/As <sub>tot</sub> (%) | B (µg/L)               | Ba (µg/L)             | Cd (µg/L)              | Co (µg/L)               | Cu (µg/L)              | Cr (µg/L)             | Fe (µg/L)                            | Mn (µg/L)                           | Ni (µg/L)                           | Pb (µg/L)                            | Zn (µg/L) |       |
| Ground-water | P1                   | 2.96      | 203                           | 0                           | 100                    | 24.3                  | 29.7                   | 0.005                   | 0.384                  | 8.08                  | 0.103                                | 0.860                               | 1.56                                | 3.56                                 | 0.037     | 12.7  |
|              | P2                   | 7.74      | 0.714                         | 0                           | 100                    | 30.0                  | 37.0                   | 0.004                   | 0.188                  | 1.72                  | 1.16                                 | 5.26                                | 0.314                               | 0.125                                | 0.034     | 3.61  |
|              | P3                   | 3.96      | 2.54                          | 100                         | 0                      | 17.7                  | 2.98                   | 0.013                   | 0.179                  | 1.13                  | 0.041                                | 4.09                                | 2.36                                | 0.131                                | 0.013     | 0.675 |
|              | P4                   | 4.53      | 190                           | n.d                         | n.d                    | 14.1                  | 86.6                   | 0.004                   | 9.79                   | 1.97                  | 0.037                                | 5.60                                | 956                                 | 6.15                                 | 0.021     | 2.41  |
|              | P5                   | 86.1      | 0.130                         | n.d                         | n.d                    | 11.0                  | 33.2                   | 0.002                   | 0.048                  | 0.326                 | 0.332                                | 40.9                                | 2.26                                | 0.783                                | 0.023     | 2.19  |
|              | P6                   | 74.2      | 0.086                         | n.d                         | n.d                    | 11.4                  | 32.6                   | 0.001                   | 0.070                  | 1.23                  | 0.182                                | 40.6                                | 0.546                               | 0.655                                | 0.041     | 9.85  |
|              | P7                   | 34.6      | 0.253                         | n.d                         | n.d                    | 13.7                  | 50.0                   | 0.004                   | 0.111                  | 0.763                 | 0.125                                | 17.3                                | 16.7                                | 0.766                                | 0.042     | 1.41  |
|              | P8                   | 2.61      | 15.8                          | n.d                         | n.d                    | 20.3                  | 5.39                   | 0.007                   | 0.041                  | 0.298                 | 0.174                                | 0.547                               | 0.152                               | 0.361                                | 0.027     | 0.624 |

Table 2 (continued)

| Sample name | Al (µg/L) | As (µg/L) | As(III)/As <sub>tot</sub> (%) | As(V)/As <sub>tot</sub> (%) | B (µg/L) | Ba (µg/L) | Cd (µg/L) | Co (µg/L) | Cu (µg/L) | Cr (µg/L) | Fe (µg/L) | Mn (µg/L) | Ni (µg/L) | Pb (µg/L) | Zn (µg/L) |
|-------------|-----------|-----------|-------------------------------|-----------------------------|----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| P9          | 0.262     | 0.005     | n.d                           | n.d                         | 1.98     | 3.16      | 0.011     | 0.053     | 3.66      | 0.008     | 0.751     | 3.14      | 0.180     | 0.912     | 7.95      |
| P10         | 11.2      | 0.078     | n.d                           | n.d                         | 118      | 72.9      | 0.016     | 6.79      | 4.17      | 0.258     | 6.56      | 121       | 11.2      | 0.071     | 34.7      |
| P11         | 67.7      | 0.038     | n.d                           | n.d                         | 15.6     | 131       | 0.141     | 5.59      | 0.410     | 0.137     | 6.05      | 657       | 24.2      | 0.015     | 39.0      |
| P12         | 13.9      | 0.360     | n.d                           | n.d                         | 11.8     | 158       | 0.005     | 5.46      | 0.188     | 0.069     | 62.6      | 581       | 6.57      | 0.035     | 1.61      |
| P13         | 4.59      | 0.048     | n.d                           | n.d                         | 4.70     | 94.5      | 0.050     | 0.227     | 1.51      | 0.055     | 1.82      | 17.7      | 1.87      | 0.030     | 10.9      |
| P14         | 10.9      | 0.028     | n.d                           | n.d                         | 6.71     | 14.7      | 0.014     | 0.285     | 0.500     | 0.063     | 19.5      | 24.0      | 0.481     | 0.369     | 8.82      |
| F1          | 7.91      | 0.183     | 0                             | 100                         | 14.0     | 458       | 0.007     | 0.048     | 1.17      | 0.035     | 3.16      | 1.87      | 0.218     | 0.045     | 3.12      |
| F2          | 3.33      | 0.149     | 0                             | 100                         | 8.04     | 275       | 0.007     | 0.058     | 0.756     | 0.145     | 0.648     | 1.20      | 0.280     | 0.014     | 1.48      |
| F3          | 5.97      | 0.298     | 0                             | 100                         | 4.57     | 201       | 0.008     | 0.027     | 1.76      | 0.082     | 2.48      | 0.354     | 0.312     | 0.037     | 2.24      |
| F4          | 2.03      | 0.365     | n.d                           | n.d                         | 27.7     | 54.8      | 0.008     | 0.197     | 8.62      | 0.113     | 1.23      | 73.6      | 2.36      | 0.020     | 5.86      |
| F5          | 43.5      | 0.681     | n.d                           | n.d                         | 45.5     | 443       | 0.049     | 0.240     | 3.29      | 0.970     | 23.5      | 4.52      | 4.38      | 0.484     | 37.5      |
| F6          | 1.44      | 2.03      | 0                             | 100                         | 7.28     | 3.35      | 0.003     | 0.035     | 0.143     | 0.026     | 1.40      | 0.273     | 0.067     | 0.020     | 34.9      |
| F7          | 11.8      | 0.116     | n.d                           | n.d                         | 14.8     | 146       | 0.006     | 0.050     | 0.431     | 0.169     | 11.7      | 0.899     | 0.677     | 0.038     | 6.13      |
| F8          | 19.1      | 0.177     | n.d                           | n.d                         | 5.81     | 284       | 0.017     | 0.011     | 0.680     | 0.295     | 2.55      | 0.639     | 0.418     | 0.121     | 57.0      |
| F9          | 22.5      | 0.043     | n.d                           | n.d                         | 3.69     | 109       | 0.005     | 0.030     | 0.417     | 6.02      | 0.432     | 1.90      | 0.659     | 0.012     | 6.62      |
| F10         | 0.593     | 0.049     | n.d                           | n.d                         | 13.0     | 178       | 0.004     | 0.006     | 1.13      | 0.021     | 0.331     | 0.240     | 0.552     | 0.031     | 1.17      |
| F11         | 1.48      | 0.061     | n.d                           | n.d                         | 24.5     | 244       | 0.004     | 0.498     | 0.680     | 0.159     | 0.530     | 0.122     | 0.165     | 0.030     | 4.49      |
| F12         | 6.98      | 15.5      | n.d                           | n.d                         | 14.1     | 191       | 0.005     | 0.024     | 0.345     | 0.220     | 2.06      | 0.138     | 0.081     | 0.006     | 1.27      |
| F13         | 5.30      | 0.371     | n.d                           | n.d                         | 72.9     | 347       | 0.023     | 0.420     | 10.8      | 8.82      | 1.22      | 0.386     | 0.218     | 0.687     | 14.4      |
| F14         | 0.638     | 0.079     | n.d                           | n.d                         | 50.9     | 1344      | 0.003     | 0.016     | 0.231     | 0.035     | 0.583     | 0.558     | 0.066     | 0.061     | 2.32      |
| F15         | 3.94      | 5.07      | n.d                           | n.d                         | 16.5     | 45.3      | 0.002     | 0.087     | 0.235     | 0.903     | 1.45      | 0.089     | 0.379     | 0.034     | 0.564     |
| F16         | 2.23      | 0.209     | n.d                           | n.d                         | 21.6     | 296       | 0.006     | 0.018     | 0.343     | 0.189     | 0.395     | 0.998     | 0.349     | 0.032     | 6.38      |
| F17         | 1.82      | 2.12      | n.d                           | n.d                         | 11.8     | 135       | 0.009     | 0.087     | 2.79      | 0.034     | 0.846     | 24.5      | 0.141     | 0.052     | 6.54      |
| F18         | 2.23      | 0.293     | n.d                           | n.d                         | 7.84     | 60.0      | 0.007     | 0.295     | 1.30      | 0.187     | 0.294     | 0.227     | 0.133     | 0.032     | 2.02      |
| F19         | 1.85      | 0.067     | n.d                           | n.d                         | 7.65     | 10.7      | 0.003     | 0.029     | 0.276     | 0.058     | 0.363     | 0.517     | 0.156     | 0.056     | 13.8      |
| F20         | 2.59      | 0.071     | n.d                           | n.d                         | 1.377    | 30.6      | 0.003     | 0.016     | 0.119     | 0.008     | 7.28      | 70.1      | 0.045     | 0.006     | 0.866     |
| F21         | 20.9      | 0.018     | n.d                           | n.d                         | 5.35     | 29.1      | 0.069     | 2.08      | 3.70      | 0.038     | 15.1      | 42.4      | 4.54      | 0.155     | 9.68      |
| F22         | 1.20      | 1.10      | n.d                           | n.d                         | 93.8     | 43.8      | 0.001     | 3.48      | 0.228     | 0.013     | 12.1      | 758       | 4.78      | 0.005     | 39.6      |
| F23         | 4.48      | 0.063     | n.d                           | n.d                         | 2.10     | 1.69      | <0.001    | 18.0      | 0.157     | 0.027     | 103       | 420       | 9.13      | 0.020     | 12.8      |
| F24         | 3.30      | 0.130     | n.d                           | n.d                         | 1.092    | 27.7      | 0.003     | 0.016     | 0.196     | 0.009     | 2.96      | 49.0      | 0.357     | 0.029     | 94.7      |
| F25         | 2.55      | 0.054     | n.d                           | n.d                         | 12.6     | 108       | 0.004     | 0.008     | 8.90      | 0.015     | 1.57      | 0.872     | 0.264     | 0.991     | 14.5      |
| F26         | 2.44      | 0.091     | n.d                           | n.d                         | 191      | 34.9      | 0.000     | 0.059     | 0.104     | 0.016     | 0.947     | 116       | 0.196     | 0.009     | 1.65      |
| F27         | 1.73      | 0.153     | n.d                           | n.d                         | 41.3     | 227       | 0.053     | 3.45      | 2.36      | 0.063     | 27.2      | 611       | 2.81      | 0.389     | 21.5      |
| F28         | 4.03      | 0.042     | n.d                           | n.d                         | 5.14     | 31.5      | 0.017     | 0.193     | 150       | 0.677     | 0.819     | 3.70      | 1.14      | 3.16      | 71.7      |
| F29         | 3.50      | 0.028     | n.d                           | n.d                         | 446      | 128       | 0.001     | 0.057     | 0.532     | 0.015     | 101       | 17.8      | 0.095     | 0.068     | 6.84      |

**Table 2** (continued)



| Sample name           | Latitude              | Longitude | Sampling date (wet (W)/dry (D) period) | Watershed               | Geological environment | Structure             | Water use              | Aquifer type            | Geological formation   | Depth (m)             |                                      |                                     |                                     |                                      |
|-----------------------|-----------------------|-----------|--|-------------------------|------------------------|-----------------------|------------------------|-------------------------|------------------------|-----------------------|--------------------------------------|-------------------------------------|-------------------------------------|--------------------------------------|
| Surface water         |                       |           |  |                         |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |
| Sirb 1                | 13.76947              | 1.65249   | 18/04/2019, D                          | Sirba                   | ES                     | River                 |                        | Sirba                   | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Sirb 2                | 13.75928              | 1.62998   | 16/05/2019, D                          | Sirba                   | ES                     | River                 |                        | Sirba                   | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Sirb 3                | 13.73168              | 1.59957   | 16/05/2019, D                          | Sirba                   | ES                     | River                 |                        | Sirba                   | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Sirb 4                | 13.66488              | 1.49981   | 16/05/2019, D                          | Sirba                   | ES                     | River                 |                        | Sirba                   | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Sirb 5                | 13.61660              | 1.45443   | 16/05/2019, D                          | Sirba                   | ES                     | River                 |                        | Sirba                   | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Sirb 6                | 13.49008              | 1.26186   | 17/07/2019, W                          | Sirba                   | ES                     | River                 |                        | Sirba                   | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Sirb 7                | 13.58827              | 1.35160   | 17/07/2019, W                          | Sirba                   | ES                     | River                 |                        | Sirba                   | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Darg 1                | 14.01499              | 0.74457   | 18/04/2019, D                          | Dargol                  | ES                     | River                 |                        | Dargol                  | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Darg 2                | 13.95177              | 0.97476   | 25/04/2019, D                          | Dargol                  | ES                     | River                 |                        | Dargol                  | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Darg 3                | 13.85238              | 1.46900   | 07/05/2019, D                          | Dargol                  | ES                     | River                 |                        | Dargol                  | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Darg 4                | 13.95177              | 0.97476   | 07/05/2019, D                          | Dargol                  | ES                     | River                 |                        | Dargol                  | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Goro 1                | 13.11141              | 1.80470   | 07/05/2019, D                          | Goroubi                 | ES                     | River                 |                        | Goroubi                 | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Goro 2                | 13.22084              | 1.86547   | 07/05/2019, D                          | Goroubi                 | ES                     | River                 |                        | Goroubi                 | LG                     | 0                     |                                      |                                     |                                     |                                      |
| Reference values      |                       |           |  |                         |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |
| CMA NN                |                       |           |  |                         |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |
| WHO guidelines        |                       |           |  |                         |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |
| CMA NF                |                       |           |  |                         |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |
| CMA NC                |                       |           |  |                         |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |
| Niger river*          |                       |           |  |                         |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |
| World river average** |                       |           |  |                         |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |
| Sample name           | Conduc-tivity (µS/cm) | T°C       | pH                                     | Ca <sup>2+</sup> (mg/L) | Cl <sup>-</sup> (mg/L) | K <sup>+</sup> (mg/L) | Na <sup>+</sup> (mg/L) | Mg <sup>2+</sup> (mg/L) | Br <sup>-</sup> (mg/L) | F <sup>-</sup> (mg/L) | HCO <sub>3</sub> <sup>-</sup> (mg/L) | NO <sub>3</sub> <sup>-</sup> (mg/L) | NO <sub>2</sub> <sup>-</sup> (mg/L) | SO <sub>4</sub> <sup>2-</sup> (mg/L) |
| Surface water         | 107                   | 32        | 6.28                                   | 5.57                    | 1.43                   | 3.47                  | 5.88                   | 2.69                    | <0.02                  | 0.17                  | 55.5                                 | <0.04                               | <0.08                               | 0.39                                 |
| Sirb 2                | 145                   | 31        | 6.20                                   | 12.0                    | 4.05                   | 5.40                  | 9.53                   | 4.75                    | <0.02                  | 0.21                  | 93.5                                 | 0.05                                | <0.08                               | 1.05                                 |
| Sirb 3                | 488                   | 31        | 6.74                                   | 21.0                    | 74.0                   | 15.4                  | 83.6                   | 11.2                    | 0.45                   | 0.58                  | 99.6                                 | 6.67                                | <0.08                               | 29.2                                 |
| Sirb 4                | 389                   | 30        | 7.24                                   | 27.1                    | 11.7                   | 25.0                  | 18.7                   | 16.6                    | 0.15                   | 0.52                  | 266                                  | 0.07                                | <0.08                               | 1.34                                 |
| Sirb 5                | 284                   | 33        | 6.43                                   | 25.5                    | 8.09                   | 10.8                  | 13.7                   | 11.0                    | <0.02                  | 0.36                  | 192                                  | <0.04                               | <0.08                               | 2.02                                 |
| Sirb 6                | 41                    | 25        | 5.80                                   | 3.02                    | 1.42                   | 3.82                  | 1.92                   | 1.03                    | <0.02                  | <0.08                 | 18.2                                 | 2.59                                | <0.08                               | 1.12                                 |
| Sirb 7                | 104                   | 25        | n.d                                    | 2.26                    | 1.14                   | 0.60                  | 1.33                   | 0.50                    | <0.02                  | <0.08                 | n.d                                  | 0.08                                | <0.08                               | 0.78                                 |
| Darg 1                | 163                   | 25        | 6.27                                   | 15.0                    | 3.24                   | 7.66                  | 7.14                   | 5.77                    | <0.02                  | 0.60                  | 104                                  | <0.04                               | <0.08                               | 3.59                                 |
| Darg 2                | 179                   | 25        | 6.21                                   | 10.5                    | 10.8                   | 12.3                  | 13.0                   | 3.90                    | 0.07                   | 0.34                  | n.d                                  | 4.68                                | <0.08                               | 8.35                                 |
| Darg 3                | 308                   | 33        | 7.24                                   | 30.6                    | 6.80                   | 7.42                  | 22.9                   | 6.14                    | <0.02                  | 0.51                  | 88.4                                 | 1.93                                | <0.08                               | 3.95                                 |
| Darg 4                | 159                   | 36        | 6.16                                   | 11.1                    | 3.96                   | 8.96                  | 12.6                   | 3.72                    | <0.02                  | 0.26                  | 89.4                                 | 4.70                                | <0.08                               | 3.15                                 |
| Goro 1                | 45                    | 25        | 5.86                                   | 3.66                    | 1.20                   | 5.22                  | 1.43                   | 1.26                    | <0.02                  | <0.08                 | 23.0                                 | 1.98                                | <0.08                               | 1.10                                 |
| Goro 2                | 30                    | 25        | 5.74                                   | 2.72                    | 0.96                   | 2.95                  | 1.00                   | 0.92                    | <0.02                  | <0.08                 | 13.9                                 | 3.06                                | 0.13                                | 0.88                                 |

**Table 2** (continued)

| Sample name             | Conductivity (µS/cm) | T°C       | pH                            | Ca <sup>2+</sup> (mg/L)     | Cl <sup>-</sup> (mg/L) | K <sup>+</sup> (mg/L) | Na <sup>+</sup> (mg/L) | Mg <sup>2+</sup> (mg/L) | Br <sup>-</sup> (mg/L) | F <sup>-</sup> (mg/L) | HCO <sub>3</sub> <sup>-</sup> (mg/L) | NO <sub>3</sub> <sup>-</sup> (mg/L) | NO <sub>2</sub> <sup>-</sup> (mg/L) | SO <sub>4</sub> <sup>2-</sup> (mg/L) |           |
|-------------------------|----------------------|-----------|-------------------------------|-----------------------------|------------------------|-----------------------|------------------------|-------------------------|------------------------|-----------------------|--------------------------------------|-------------------------------------|-------------------------------------|--------------------------------------|-----------|
| <b>Reference values</b> |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| CMA NN                  |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| WHO guidelines          |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| CMA NF                  |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| CMA NC                  |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| Niger river*            |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| World river average*    |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| Sample name             | Al (µg/L)            | As (µg/L) | As(III)/As <sub>tot</sub> (%) | As(V)/As <sub>tot</sub> (%) | B (µg/L)               | Ba (µg/L)             | Cd (µg/L)              | Co (µg/L)               | Cu (µg/L)              | Cr (µg/L)             | Fe (µg/L)                            | Mn (µg/L)                           | Ni (µg/L)                           | Pb (µg/L)                            | Zn (µg/L) |
| Surface water           |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| Sirb 1                  | 7.52                 | 0.738     | 100                           | 0                           | 7.71                   | 50.8                  | 0.210                  | 0.081                   | 1.72                   | 0.104                 | 9.02                                 | 2.30                                | 0.786                               | 0.096                                | 4.30      |
| Sirb 2                  | 20.2                 | 1.72      | 100                           | 0                           | 11.3                   | 109                   | 0.013                  | 0.135                   | 1.35                   | 0.082                 | 21.3                                 | 9.88                                | 0.902                               | 0.034                                | 4.97      |
| Sirb 3                  | 28.0                 | 2.91      | 70                            | 30                          | 38.9                   | 280                   | 0.041                  | 0.865                   | 5.67                   | 0.300                 | 21.3                                 | 4.36                                | 3.77                                | 0.060                                | 8.80      |
| Sirb 4                  | 17.9                 | 2.25      | 100                           | 0                           | 32.1                   | 660                   | 0.035                  | 0.455                   | 2.20                   | 0.108                 | 13.4                                 | 37.1                                | 1.96                                | 0.037                                | 11.0      |
| Sirb 5                  | 15.6                 | 1.65      | 100                           | 0                           | 22.8                   | 256                   | 0.461                  | 0.441                   | 2.37                   | 0.098                 | 5.84                                 | 68.9                                | 1.32                                | 0.046                                | 12.0      |
| Sirb 6                  | 821                  | 0.614     | n.d                           | n.d                         | 12.7                   | 62.7                  | 0.052                  | 0.437                   | 3.45                   | 1.08                  | 590                                  | 21.7                                | 1.56                                | 0.521                                | 15.7      |
| Sirb 7                  | 133                  | 0.531     | n.d                           | n.d                         | 11.9                   | 55.7                  | 0.033                  | 1.99                    | 1.06                   | 0.196                 | 82.5                                 | 66.7                                | 1.11                                | 0.112                                | 10.2      |
| Darg 1                  | 563                  | 0.576     | 0                             | 100                         | 26.0                   | 141                   | 0.115                  | 0.372                   | 3.00                   | 0.965                 | 431                                  | 12.7                                | 1.49                                | 0.520                                | 3.25      |
| Darg 2                  | 550                  | 0.306     | n.d                           | n.d                         | 74.4                   | 296                   | 0.003                  | 0.190                   | 3.29                   | 1.00                  | 234                                  | 1.01                                | 1.39                                | 0.146                                | 3.07      |
| Darg 3                  | 11.2                 | 0.339     | n.d                           | n.d                         | 18.3                   | 272                   | 0.052                  | 0.133                   | 1.29                   | 0.133                 | 15.2                                 | 1.39                                | 0.741                               | 0.042                                | 4.99      |
| Darg 4                  | 356                  | 0.153     | n.d                           | n.d                         | 15.9                   | 179                   | 0.026                  | 0.404                   | 1.91                   | 0.306                 | 181                                  | 161                                 | 0.694                               | 0.363                                | 3.64      |
| Goro 1                  | 135                  | 0.186     | n.d                           | n.d                         | 13.8                   | 39.8                  | 0.028                  | 0.271                   | 2.16                   | 0.298                 | 86.8                                 | 8.69                                | 1.16                                | 0.070                                | 6.77      |
| Goro 2                  | 19.9                 | 0.130     | n.d                           | n.d                         | 7.37                   | 27.2                  | 0.020                  | 0.140                   | 1.13                   | 0.164                 | 15.4                                 | 3.37                                | 0.715                               | 0.037                                | 5.46      |
| Reference values        |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| CMA NN                  | 200                  | 10        |                               |                             | 2400                   | 1300                  | 5                      | 1000                    | 1000                   | 50                    | 50                                   | 100                                 | 70                                  | 50                                   | 5000      |
| WHO                     | 200                  | 10        |                               |                             | 2400                   | 1300                  | 3                      | 2000                    | 2000                   | 50                    | 50                                   | 400                                 | 70                                  | 10                                   | 3000      |
| guide-lines             |                      |           |                               |                             |                        |                       |                        |                         |                        |                       |                                      |                                     |                                     |                                      |           |
| CMA NF                  | 200                  | 10        |                               |                             | 1000                   | 700                   | 5                      | 2000                    | 2000                   | 50                    | 50                                   | 50                                  | 20                                  | 10                                   |           |
| CMA NC                  | 10                   | 10        |                               |                             | 5000                   | 2000                  | 7                      | 2000                    | 2000                   | 50                    | 50                                   | 120                                 |                                     | 5                                    |           |
| Niger river*            | 76                   |           |                               |                             | 3.2                    | 30                    |                        | 0.040                   | 0.63                   | 0.45                  | 105                                  | 0.50                                | 0.29                                | 0.039                                | 0.89      |
| World river average*    | 32                   | 0.62      |                               |                             | 10.2                   | 23                    | 0.08                   | 0.148                   | 1.48                   | 0.7                   | 66                                   | 34                                  | 0.801                               | 0.079                                | 0.60      |

\*From Gaillardet et al., 2003  
n.d. not determined

**Fig. 3** Comparison of dissolved concentration ( $\mu\text{g/L}$ ) of trace elements in the two contrasted geological environments CT (in red) and LG (in blue)



**Fig. 4** Distribution of As concentrations in the studied area






Fig. 5 Distribution of Mn concentrations in the studied area

in SPM and more extensively for B ( $EF_{\text{average}} = 6 \pm 4$ ) and Cr ( $EF_{\text{average}} = 8 \pm 5$ ) in sediments.

## Discussion

### Arsenic in groundwaters and rivers

Our study evidenced 23% of groundwater samples that exceeded drinking water guidelines for As or Mn. For As, exceedances of guidelines were exclusively encountered in the LG geologic formation, in alterite greenstone aquifers of the Birimian formations. There was no boreholes or wells with high As concentrations in the alluvial aquifers of the Continental Terminal. The four groundwater samples with the highest arsenic concentrations have been taken from wells or boreholes located in or near the gold artisanal mining sites of Mbanga ([As] = 203 µg/L), Boulounjounga ([As] = 190 µg/L) and Komabangou ([As] = 15.8 µg/L and 15.5 µg/L). The wells from Mbanga and Boulounjounga have been dug to exploit vein gold deposits, and reached

the water table, thus providing a water resource to the surrounding population. At Komabangou, the two boreholes exceeding 15 µg/L have been drilled for drinking water supply. Previous data from Tankari et al. (2014, 2019) showed similar values in groundwater at Komabangou (As = 19 µg/L, Tankari et al. 2014; As = 3 µg/L, Tankari et al. 2019) and in water from the gravimetric gold extraction process (As = 20 µg/L) while the As concentration was hundred times higher (As = 2429 µg/L) in the water from the cyanidation gold extraction process. In soils, average As concentrations were two orders of magnitude higher in the gold exploitation zone than outside this zone (Tankari et al. 2019). Thus, Au mineralisation was a source of high-As groundwater at Komabangou. Our study revealed that, besides Komabangou, two other artisanal gold mining sites (Mbanga and Boulounjounga) in the Nigerien Liptako exhibited even higher groundwater As concentrations. The reason for such elevated As concentrations in wells P1 (203 µg/L) and P4 (190 µg/L) may be related to their original purpose (gold extraction rather than drinking water supply) and age (they have been exploited since the 1980's). It

**Table 3** Concentrations and enrichment factors in SPM and sediments from the rivers Sirba, Dargol and Goroubi. Al and Fe concentrations in %, other trace element concentrations in µg/g. World river averages are given for SPM when available (Viers et al. 2009)

| River          | Sample name | Al          |          | As          |          | B    |          | Ba   |          | Cd          |            | Co     |          | Cr          |             |       |       |      |     |             |     |      |     |      |     |              |            |  |
|----------------|-------------|-------------|----------|-------------|----------|------|----------|------|----------|-------------|------------|--------|----------|-------------|-------------|-------|-------|------|-----|-------------|-----|------|-----|------|-----|--------------|------------|--|
|                |             | SPM         | Sediment | SPM         | Sediment | SPM  | Sediment | SPM  | Sediment | SPM         | Sediment   | SPM    | Sediment | SPM         | Sediment    |       |       |      |     |             |     |      |     |      |     |              |            |  |
|                |             | %           | %        | µg/g        | EF       | µg/g | EF       | µg/g | EF       | µg/g        | EF         | µg/g   | EF       | µg/g        | EF          | µg/g  | EF    |      |     |             |     |      |     |      |     |              |            |  |
| Sirba          | Sirb 1      | 11.2        | 1.9      | 12.8        | 6.1      | 4.2  | 11.9     | 32.1 | 1.5      | 15.7        | 4.5        | 499    | 0.7      | 144.5       | 1.1         | 21.0  | 153.4 | 0.03 | 1.4 | 24.9        | 1.8 | 5.6  | 2.4 | 123  | 2.5 | 42.1         | 5.1        |  |
|                | Sirb 2      | 0.95        | 0.8      | 1.80        | 10.1     | 3.7  | 26.4     | 2.12 | 1.2      | 17.6        | 12.5       | 126    | 1.9      | 71.9        | 1.4         | 0.035 | 3.0   | 0.03 | 3.2 | 3.27        | 2.8 | 3.8  | 4.0 | 11.9 | 2.9 | 49.8         | 15.2       |  |
|                | Sirb 3      | 4.65        | 2.1      | 5.19        | 6.0      | 22.5 | 58.7     | 5.94 | 0.7      | 22.5        | 5.9        | 124    | 0.4      | 188.2       | 1.3         | 0.027 | 0.5   | 0.05 | 1.8 | 9.60        | 1.7 | 17.1 | 6.7 | 65.9 | 3.3 | 188.5        | 21.1       |  |
|                | Sirb 4      | 2.41        | 1.6      | 2.95        | 6.5      | 5.0  | 17.2     | 4.27 | 1.0      | 18.6        | 6.4        | 140    | 0.8      | 437.8       | 4.1         | 0.020 | 0.7   | 0.03 | 1.5 | 5.34        | 1.8 | 6.5  | 3.4 | 32.3 | 3.1 | 42.5         | 6.2        |  |
|                | Sirb 5      | 1.28        | 0.9      | 1.73        | 7.3      | 1.8  | 10.1     | 4.58 | 1.9      | 7.9         | 4.6        | 69.9   | 0.8      | 157.5       | 2.5         | 0.027 | 1.7   | 0.02 | 1.8 | 2.85        | 1.8 | 2.5  | 2.2 | 15.1 | 2.7 | 28.9         | 7.1        |  |
|                | Sirb 6      | 1.38        | 0.4      | 2.10        | 8.2      | 1.8  | 25.2     | 1.86 | 0.7      | 9.1         | 12.4       | 31.6   | 0.3      | 27.4        | 1.0         | 0.020 | 1.2   | 0.01 | 1.5 | 1.92        | 1.1 | 1.6  | 3.3 | 13.7 | 2.3 | 15.4         | 9.1        |  |
|                | Sirb 7      | 6.52        | 2.7      | 17.0        | 14.0     | 6.2  | 12.1     | 30.3 | 2.5      | 28.0        | 5.5        | 412    | 0.9      | 272.3       | 1.5         | 0.067 | 0.8   | 0.04 | 1.3 | 20.8        | 2.6 | 12.2 | 3.6 | 118  | 4.1 | 68.3         | 5.7        |  |
| Dargol         | Darg 1      | 11.3        | 5.6      | 5.63        | 2.7      | 5.2  | 4.9      | 14.7 | 0.7      | 13.3        | 1.3        | 237    | 0.3      | 253.4       | 0.7         | 0.268 | 2.0   | 0.06 | 0.9 | 11.9        | 0.8 | 14.1 | 2.0 | 104  | 2.1 | 103.3        | 4.2        |  |
|                | Darg 2      | 10.7        | 2.1      | 5.28        | 2.6      | 1.7  | 4.3      | 11.4 | 0.6      | 10.3        | 2.6        | 401    | 0.5      | 165.9       | 1.1         | 0.057 | 0.4   | 0.02 | 0.7 | 18.3        | 1.4 | 5.8  | 2.2 | 138  | 3.0 | 40.6         | 4.4        |  |
| Goroubi        | Goro 1      | 5.22        | 1.7      | 1.95        | 2.0      | 1.4  | 4.4      | 4.77 | 0.5      | 9.6         | 3.0        | 125    | 0.4      | 144.6       | 1.2         | 0.018 | 0.3   | 0.02 | 0.8 | 6.52        | 1.0 | 5.7  | 2.7 | 53.6 | 2.4 | 37.4         | 5.0        |  |
|                | Goro 2      | 11.6        | 3.4      | 6.20        | 2.9      | 7.1  | 11.1     | 13.4 | 0.6      | 15.8        | 2.5        | 288    | 0.4      | 163.3       | 0.7         | 0.044 | 0.3   | 0.06 | 1.4 | 19.7        | 1.4 | 11.4 | 2.7 | 159  | 3.1 | 84.2         | 5.6        |  |
| World River av |             | <b>8.72</b> |          | <b>36.3</b> |          |      |          |      |          |             | <b>522</b> |        |          |             | <b>1.55</b> |       |       |      |     | <b>22.5</b> |     |      |     |      |     |              | <b>130</b> |  |
|                |             |             |          |             |          |      |          |      | No data  |             |            |        |          |             |             |       |       |      |     |             |     |      |     |      |     |              |            |  |
| River          | Sample name | Cu          |          | Fe          |          | Mn   |          | Ni   |          | Pb          |            | Zn     |          |             |             |       |       |      |     |             |     |      |     |      |     |              |            |  |
|                |             | SPM         | Sediment | SPM         | Sediment | SPM  | Sediment | SPM  | Sediment | SPM         | Sediment   | SPM    | Sediment |             |             |       |       |      |     |             |     |      |     |      |     |              |            |  |
|                |             | µg/g        | EF       | %           | EF       | %    | EF       | µg/g | EF       | µg/g        | EF         | µg/g   | EF       |             |             |       |       |      |     |             |     |      |     |      |     |              |            |  |
| Sirba          | Sirb 1      | 52.6        | 1.5      | 11.0        | 1.9      | 5.26 | 1.1      | 1.4  | 1.7      | 2832        | 3.4        | 214.6  | 1.5      | 71.5        | 2.6         | 13.3  | 2.8   | 22.8 | 0.8 | 6.7         | 1.4 | 137  | 1.4 | 18.9 | 1.1 |              |            |  |
|                | Sirb 2      | 5.03        | 1.7      | 5.8         | 2.5      | 0.49 | 1.2      | 1.0  | 3.0      | 816         | 11.5       | 285.8  | 5.1      | 6.38        | 2.7         | 6.2   | 3.3   | 2.80 | 1.2 | 8.5         | 4.6 | 12.1 | 1.4 | 12.4 | 1.9 |              |            |  |
|                | Sirb 3      | 21.5        | 1.5      | 37.1        | 5.8      | 2.11 | 1.0      | 8.2  | 9.2      | 242         | 0.7        | 407.1  | 2.7      | 29.2        | 2.5         | 32.8  | 6.4   | 5.82 | 0.5 | 8.5         | 1.7 | 29.8 | 0.7 | 35.6 | 2.0 |              |            |  |
|                | Sirb 4      | 12.0        | 1.6      | 12.8        | 2.6      | 1.21 | 1.2      | 1.5  | 2.1      | 468         | 2.6        | 1292.4 | 11.1     | 15.4        | 2.6         | 12.2  | 3.1   | 3.36 | 0.6 | 5.7         | 1.5 | 18.4 | 0.9 | 16.5 | 1.2 |              |            |  |
|                | Sirb 5      | 5.77        | 1.5      | 5.4         | 1.9      | 0.61 | 1.1      | 0.7  | 1.8      | 373         | 3.9        | 146.1  | 2.1      | 8.12        | 2.6         | 4.7   | 2.0   | 2.08 | 0.7 | 4.3         | 1.8 | 11.1 | 1.0 | 8.9  | 1.1 |              |            |  |
|                | Sirb 6      | 5.12        | 1.2      | 3.6         | 2.9      | 0.69 | 1.2      | 0.5  | 3.2      | 36.5        | 0.4        | 58.4   | 2.0      | 7.05        | 2.1         | 3.5   | 3.6   | 1.85 | 0.5 | 2.2         | 2.2 | 11.2 | 0.9 | 6.0  | 1.7 |              |            |  |
|                | Sirb 7      | 40.6        | 2.0      | 18.9        | 2.2      | 4.64 | 1.6      | 2.7  | 2.2      | 622         | 1.3        | 399.0  | 2.0      | 55.7        | 3.4         | 23.9  | 3.5   | 14.5 | 0.9 | 9.7         | 1.4 | 65.2 | 1.1 | 30.1 | 1.2 |              |            |  |
| Dargol         | Darg 1      | 30.2        | 0.9      | 33.9        | 1.9      | 3.93 | 0.8      | 3.7  | 1.5      | 220         | 0.3        | 795.0  | 1.9      | 55.0        | 2.0         | 48.1  | 3.5   | 12.4 | 0.4 | 15.2        | 1.1 | 73.1 | 0.7 | 66.7 | 1.4 |              |            |  |
|                | Darg 2      | 40.5        | 1.2      | 11.8        | 1.8      | 5.16 | 1.1      | 1.3  | 1.4      | 494         | 0.6        | 210.4  | 1.3      | 68.1        | 2.6         | 16.6  | 3.1   | 17.4 | 0.7 | 6.5         | 1.2 | 74.2 | 0.8 | 23.1 | 1.2 |              |            |  |
| Goroubi        | Goro 1      | 15.9        | 1.0      | 8.1         | 1.5      | 1.87 | 0.8      | 1.0  | 1.4      | 117         | 0.3        | 144.3  | 1.1      | 29.6        | 2.3         | 13.5  | 3.2   | 6.35 | 0.5 | 5.8         | 1.4 | 36.0 | 0.8 | 19.0 | 1.3 |              |            |  |
|                | Goro 2      | 39.5        | 1.1      | 16.6        | 1.6      | 5.31 | 1.0      | 5.5  | 3.7      | 268         | 0.3        | 261.4  | 1.0      | 78.2        | 2.7         | 25.3  | 3.0   | 21.9 | 0.8 | 15.6        | 1.8 | 101  | 1.0 | 56.7 | 1.9 |              |            |  |
| World River av |             | <b>75.9</b> |          | <b>5.81</b> |          |      |          |      |          | <b>1679</b> |            |        |          | <b>74.5</b> |             |       |       |      |     |             |     |      |     |      |     | <b>208.0</b> |            |  |

n.d. not determined

can be hypothesized that digging for decades toward deeper and deeper horizons in the mineralized zone favoured water/rock interaction and subsequent As mobilization. The process could take a certain amount of time; the contamination could not be effective until months or years after the beginning of pumping (Bonnemaison 2005). In this respect, well P3, located in Sefa Moussa, a few kilometres from well P1, had lower As concentration ( $As = 2,5 \mu\text{g/L}$ ). Gold extraction began in 2017 at this site. Similarly, well P5, located a few kilometres from well P4, had lower As concentrations ( $As = 0.13 \mu\text{g/L}$ ). This well was dug especially for drinking water supply; thus, groundwater at this site was not readily to interact directly with gold mineralization.

The detection of high-As groundwater at several sites in the study area demonstrates that the problem could be widely distributed in the gold extraction zones in Nigerien Liptako. These results agree with those of Smedley (1996) and Smedley et al. (2007) who reported groundwater As concentrations up to  $141 \mu\text{g/L}$  in similar Birimian lithology in northern Ghana and up to  $1630 \mu\text{g/L}$  in the region of Ouahigouya in Burkina Faso. In all these regions, a high spatial variability in groundwater As concentrations was evidenced at a local scale, which makes a prediction of groundwater As concentrations difficult. Regarding the primary arsenic source, arsenic-rich pyrite and arsenopyrite have been associated to gold mineralisation in the Birimian greenstone belt (e.g. Milési et al. 1989, 1992; Bourges et al. 1998; Smedley et al. 2007; Béziat et al. 2008; Sangaré et al. 2014). The oxidation of sulphide minerals is an important As mobilization mechanism (Smedley et al. 2002). Here, bulk analysis of the rocks collected in the vicinity of the artisanal mining sites showed As concentrations slightly higher than common values in shale and diorite. SEM–EDS analyses showed the presence of sulphide minerals, but no detectable levels of As in these sulphides. The only As source identified by SEM was related to Fe-oxides. Sample of schist showed Fe-oxides with 0.8 wt.% As. Secondary Fe-oxides have been suggested as a possible primary source of arsenic in groundwater in zones of mineralised Birimian volcano-sedimentary rocks in Burkina Faso (Smedley et al. 2007). Sulphate released during the initial oxidation of sulphides may have been flushed from the aquifer some time ago. Arsenic may be released from Fe-oxides by desorption. This process is favoured at high pH due to the anionic character of arsenate that competes for sorption sites with bicarbonate at the surface of Fe-oxides (Smedley and Kinniburgh 2002). Results of arsenic speciation in the present study showed that arsenate predominated in the well from Mbanga where the highest As concentration ( $203 \mu\text{g/L}$ ) has been measured. However, no correlations were evidenced in the present study between pH or bicarbonate concentrations and arsenic concentrations, probably because of the few high As concentration data available.

The Boulounjouna and Mbanga artisanal gold mining sites where the two highest groundwater As concentrations have been evidenced are both located near the Sirba River. In this river, gold miners extract gold by panning. Our analysis showed that dissolved As concentrations in the river Sirba were higher than in the rivers Dargol and Goroubi and also higher than the world river average. SPM and sediments of the Sirba river were the most enriched with As. Moreover, station Sirb3, located downstream a gold-panning area exhibited higher dissolved As concentration than station Sirb6, located upstream. This suggests that artisanal gold extraction in the river Sirba had an impact on arsenic mobilization to surface water, although the resulting As enrichment was relatively low. The predominance of arsenite upon arsenate suggests that As(V), which is the thermodynamically stable species in the oxidizing conditions of the Sirba river remains immobilized on SPM and sediments. At stations along the river Dargol, draining the Komabangou gold extraction area, where groundwater As concentrations around  $16 \mu\text{g/L}$  have been measured, no As enrichment was evidenced in surface water. The strong correlation ( $r^2 = 0.83$ ) between As and Fe in sediments from the three rivers suggests that Fe-oxides control As mobility in surface waters in the study area, as generally observed (Smedley and Kinniburgh 2002).

### Other trace elements in groundwaters and rivers

Nine groundwater samples, corresponding to 21% of the samples, had Mn concentrations exceeding the French drinking water guidelines ( $50 \mu\text{g/L}$ ), seven exceeded the Nigerien drinking water guideline ( $100 \mu\text{g/L}$ ) and five exceeded the WHO guideline ( $400 \mu\text{g/L}$ ). Several studies have shown that Mn could have neurodevelopmental effects in children exposed to concentrations higher than  $50 \mu\text{g/L}$  (WHO Guideline 2017). The risk of neurotoxic effects on children starts to increase when the concentration in drinking water exceeds  $100 \mu\text{g/L}$  (Bouchard et al. 2011). The highest concentration of Mn ( $956 \mu\text{g/L}$ ) was found together with arsenic ( $190 \mu\text{g/L}$ ) in the well of the artisanal gold mining site of Boulounjouna. At this site, the well was close to a manganese deposit highlighted in the Niger Mineral Plan (Franconi et al. 1984). Several manganese deposits have been identified in the West African craton including in Mali, Burkina Faso, Ghana and Côte d'Ivoire with occurrences in the volcano-sedimentary belts (Hein and Tshibubudze 2016). At Komabangou, Tankari et al. (2014, 2019) found Mn concentrations of  $98 \mu\text{g/L}$  and  $36 \mu\text{g/L}$  in groundwater while in gravimetric gold processing waters and cyanidation gold processing waters, Mn concentrations reached  $1.60 \text{ mg/L}$  and  $108.43 \text{ mg/L}$ , respectively, suggesting a link between Mn and gold mineralisation. Our analysis of rocks clearly showed the presence of manganese oxides in diorite sample

from the quartz vein and in the shale sample. These Mn oxides explain the high Mn concentrations in the groundwater sample from Boulounjunga. However, as with As, the high-Mn groundwaters were very locally distributed in the right bank area. Mn enrichment in surface water from the rivers Sirba, Dargol and Goroubi, compared to the Niger River, and local Mn enrichment in SPM and sediments compared to UCC confirm the influence of Mn deposits on water geochemistry in the right bank tributaries.

The other groundwater samples with high Mn concentrations were located in the Ct<sup>3</sup> aquifer, mainly along the left bank tributaries of the Niger River. High manganese concentrations had previously been highlighted in this area; at Tillakaina village, in the Tillabery region, Salihou-Djari et al. (2018) found Mn concentration of 140 µg/L in the Ct<sup>3</sup> aquifer waters. The processes at work for Mn mobilization in this area deserve further research.

Among the other trace elements, Ba, B and Ni showed a few localized groundwater enrichments. The origin of Ba in the LG geological environment was clearly related to the gold mineralisation according to the high content (85 mg/g) in the sample of diorite from the quartz vein, while the origin of Ba, B and Ni enrichments in groundwater in the CT geological environment will deserve further research.

## Conclusion

Arsenic and manganese are contaminants of concern in groundwaters of the Nigerien Liptako. Arsenic concentrations exceeding drinking water guidelines (> 10 µg/L) were associated to alterite aquifer of the Birimian greenstone belt. The highest As contents were evidenced in old wells mined for gold extraction since the 1980s and used for drinking water purpose, suggesting a relationship between the elevated arsenic concentrations found in groundwater and gold mining. The origin of As in the water of the alterite aquifer is probably related to desorption from iron/manganese oxides. Wells and boreholes with high As concentrations were highly dispersed throughout the Nigerien Liptako and it is therefore difficult to predict the As concentration at the regional scale and at the scale of a mining site as well. In the rivers, As enrichment was evidenced in SPM and sediments draining artisanal gold mining sites but dissolved As did not exceed ~ 1 µg/L. Manganese is another trace element of concern in groundwaters of the studied area. Concentrations exceeding WHO drinking water guidelines (400 µg/L) were recorded in aquifers of the Continental Terminal geological environment.

This study showed that it is essential to carry out an inventory of the concentrations of arsenic and other trace elements in the boreholes and wells of all the former artisanal gold mining sites in Nigerien Liptako, to characterize

the polluted areas and to find alternative water supply for the population living in the vicinity of these sites. Further investigations would be necessary to assess the level of As and Mn exposure of the populations living in these areas. Because of the development of artisanal gold mining in the region and the pressure on water resource availability, the establishment of long-term monitoring programs for arsenic and trace metal levels in boreholes used as water supplies by the population in gold-rich zones is absolutely required. Furthermore, non-geologic pollutants such as mercury and cyanides, used in gold extraction, should be investigated in the water resources in the area.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s12665-022-10639-8>.

**Funding** Ministère des Affaires Etrangères.

**Data availability** The authors confirm that the data supporting the findings of this study are available within the article [and/or] its supplementary materials.

## Declarations

**Conflict of interest** The authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript.

## References

- Abdou A., Bonnot H., Bory Kadey D., Chalamet D., Saint Martin M., Younfa I., 1998. Notice explicative des cartes géologiques du liptako à 1/100 000 et 1/200 000, Ministère des Mines et de la géologie, Rep. Niger.
- Adama S, Christophe DB, Hervé H (2017) Knowledge and perceptions of health and environmental risks related to artisanal gold mining by the artisanal miners in Burkina Faso: a cross-sectional survey. *Pan Afr Med J* 2017(27):280
- Adama T, Blaise S, Pierre IG (2012) Arsenic levels in tube-wells water, food, residents' urine and the prevalence of skin lesions in Yatenga Province, Burkina Faso. *Interdiscip Toxicol* 5(1):38–41
- Ama SI, Liègeois JP, Pouclet A (1996) Evolution d'un arc insulaire océanique birimien précoce au Liptako nigérien (Sirba): géologie, géochronologie et géochimie. *J Afr Sci* 22(3):235–254
- Babaye AMS, Orban P, Ousmane B, Favreau G, Brouyère S, Al Darsargues (2018) Characterization of recharge mechanisms in a Precambrian basement aquifer in semi-arid southwest Niger. *Hydrogeol J* 27:475–491
- Barro-Traoré F, Tiendrébéogo SRM, Lallogo S, Tiendrébéogo S, Dabal M, Ouédraogo H (2008) Manifestations cutanées de l'arcénicisme au Burkina Faso: aspects épidémiologiques et cliniques. *Mali Med* 2008(1):7–11
- Béziat D, Dubois M, Debat P, Nikiéma S, Stefano S, Tollon F (2008) Gold metallogeny in the Birimian Craton of Burkina Faso (West Africa). *J Afr Earth Sc* 50:215–233
- Bintou AM, Konan EK, Oumar S (2020) Évaluation de la dynamique de l'occupation du sol engendrée par l'orpaillage dans le bassin

- de Banankoro, Région de Koulikoro Sud-Ouest du Mali. *Afrique Sci* 17(3):28–39
- Bokar H, Traoré AZ, Mariko A, Diallo T, Traoré A, Sy A, Soumaré O, Dolo A, Bamba F, Sacko M, Touré O (2020) Geogenic influence and impact of mining activities on water, soil and plants in surrounding area of Morila Mines, Mali. *J Geochem Explor* 209:106429
- Bonnemaison M (2005) L'eau, facteur de libération de l'arsenic naturel. *Géosciences* 2:54–59
- Bouchard MF, Sauvé S., Barbeau B., Legrand M., Brodeur M.-E., Bouffard T., Limoges E., Bellinger D. C., Mergler D. (2011) Intellectual impairment in school-age children exposed to manganese from drinking water. *Environ Health Perspect* 119(1):138–143. <https://doi.org/10.1289/ehp.1002321>
- Bourges F, Debat P, Tollon F, Munoz M, Ingles J (1998) The geology of Taparko gold deposit, Birimian greenstone belt, Burkina Faso. *West African Mineralium Deposita* 33:591–605
- Bretzler A, Lalanne F, Nikiema J, Podgorski J, Pfenninger N, Berg M, Schirmer M (2017) Groundwater arsenic contamination in Burkina Faso, West Africa: predicting and verifying region at risk. *Sci Total Environ* 584–585(2017):958–970
- Cheilletz A., Barbey P., Lama C., Pons J., Zimmermann J. L., Dautel, D. (1994). Age de refroidissement de la croûte juvénile Birimienne d'Afrique de l'Ouest, données U–Pb, Rb–Sr et K–Ar sur les formations à 21 Ga au SW-Niger. *Comptes Rendus de l'Académie des Sciences Paris* 319:435–442.
- Dupuis D, Pons J, Prost AE (1991) Mise en place de plutons et caractérisation de la déformation birimienne au Niger occidental. *Compte Rendus Acad. Sci. Paris* 312:769–776
- Favreau G (2000) Caractérisation et modélisation d'une nappe phréatique en hausse au Sahel. Université de Paris XI, France, Thèse de doctorat, p 258
- Favreau G, Cappelaere B, Massuel S, Leblanc M, Boucher M, Boulain N, Leduc C (2009) Land clearing, climate variability, and water resources increase in semiarid Southwest Niger: a review. *Water Resources Res* 45:W00A16
- Franconi A, Joo J, Zibo I (1984) Plan minéral de la République du Niger, Tome IV, 2ème Volume ; Etude spécifique des principales substances minérales et leur contexte géologique p. 592.
- Gaillardet J, Viers J, Dupré B (2014) Trace elements in river waters. In: *Treatise on geochemistry*, pp. 195–235.
- Greigert, J., 1966. Descriptions des formations crétacées et tertiaires du bassin des Illummeden (Afrique occidentale), Editions du B.R.G.M./Dir. Mines Géol. Rép. Niger, Paris.
- Guéro A (2003) Étude des relations hydrauliques entre les différentes nappes du complexe sédimentaire de la bordure sud-ouest du bassin des Iullemeden (Niger): approches géochimique et hydrodynamique. Thèse de doctorat, Univ. Paris XI Orsay, France, p 265
- Guiraud R., Travi Y., 1990. Synthèse des connaissances l'hydrogéologie de l'Afrique de l'Ouest. Socle cristallin et cristallophyllien et sédimentaire ancien. Univ. Avignon – CEFIGRE, 2<sup>e</sup> édi., p. 147
- Hein KAA, Tshibubudze A (2016) Manganese deposits of the Oudalan province in the northeast of Burkina Faso, and Ansongo Inlier in eastern Mali. *Ore Geol Rev* 78:645–651
- Héry M, Casiot C, Resongles E, Gallice Z, Bruneel O, Desoeuvre A, Delpoux S (2014) Release of arsenite, arsenate and methyl-arsenic species from streambed sediment affected by acid mine drainage: a microcosm study. *Environ Chem* 11(5):514–524
- Ibrahim OZ, Tankari AD, Guero Y, Idi MFM, Feidt C, Sterckeman T, Echevarria G (2019) Distribution spatiale des éléments traces métalliques dans les sols de la zone aurifère de Komabangou au Niger. *Int J Biol Chem Sci* 13(1):557–573
- Kamagaté B, Seguis L, Favreau G, Seidel JL, Descloitres M, Affaton P (2007) Processus et bilan des flux hydriques d'un bassin versant de milieu tropical de socle au Bénin (Donga, haut Ouémé). *C r Geoscience* 339:418–429
- Leduc C, Karbo A (1996) Variabilité spatio-temporelle de l'impact de l'infiltration sur la nappe phréatique du Continental terminal (Hapex-Sahel, Niger). In : Hoepffner Michel (ed.), Lebel Thierry (ed.), Monteny Bruno (ed.). *Interactions surface continentale/atmosphère : l'expérience HAPEX-Sahel*. Paris : ORSTOM, 449–462. (Colloques et Séminaires). *Journées Hydrologiques*, 10., Montpellier (FRA), 1994/09/13–14. ISBN 2–7099–1307–0.
- Leduc C, Taupin JD (1997) Hydrochimie et recharge de la nappe phréatique du Continental Terminal (Niamey, Niger). *Hydrochemistry* 244:235–243
- Machens E (1973) Contribution à l'étude des formations du socle cristallin et de la couverture sédimentaire de l'ouest de la république du Niger Ministère des Mines, de la Géologie et de l'Hydraulique de la république du Niger. *Direction des mines et de la géologie* 4(82):167
- Milési JP, Feybesse JL, Ledru P, Dommanget A, Ouedraogo MF, Marcoux E, Prost AE, Vinchon C, Sylvain JP, Johan V, Tegye M, Calvez JY, Lagny P (1989) Les minéralisations aurifères de l'Afrique de l'Ouest. *Chronique De La Recherche Minière* 497:3–98
- Milési JP, Ledru P, Feybesse JL, Dommanget A, Marcoux E (1992) Early Proterozoic ore deposits and tectonics of the Birimian orogenic belt West Africa. *Precambr Res* 58(1–4):305–344
- Ochieng GM, Seanego ES, Nkwonta OI (2010) Impacts of mining on water resources in South Africa: a review. *Sci Res Essays* 5(22):3351–3357
- Ouedraogo O, Amyot M (2013) Mercury, arsenic and selenium concentrations in water and fish from sub-Saharan semi-arid freshwater reservoirs (Burkina Faso). *Sci Total Environ* 444:243–254
- Ousmane B, Daddy A, Soumaila A, Margueron T, Boubacar A, Garba Z (2006) Groundwater contamination in the Niamey urban area, Niger. In: Xu Y, Usher BH (eds) *Groundwater pollution in Africa*. Taylor and Francis, Balkema, pp 169–179
- Ousmane B (1988). Étude géochimique et isotopique des aquifères du socle de la bande sahéenne du Niger (Liptako, Sud-Maradi et Zinder-est). Thèse sciences. Université A.M. de Niamey, Niger, p. 152
- Pons J, Barbey P, Dupuis D, Léger JM (1995) Mechanism of emplacement and structure of a 2.1 Ga juvenile continental crust: the Birimien of southwestern Niger. *Precambrien Res* 70:281–301
- Poulet A, Prost A, E, Ama S. I., Lapiere H., (1990) Les ceintures birimiennes du Niger occidental (Protérozoïque inférieur), nouvelles données pétrologiques et structurales des formations métavolcaniques. *Comptes Rendus de l'Académie des Sciences, Paris. Earth Sci* 311:333–340
- Salihou-Djari MM, Stoleriu CC, Saley MB, Mihu-Pintilie. Alin, Romanescu G. (2018) Groundwater quality analysis in warm semi-arid climate of Sahel countries: Tillabéri region, Niger. *Carpathian J Earth Environ Sci* 13(1):277–290
- Salle LGL, C., Marlin C., Leduc C., Massault M., Favreau G., (2001) Renewal rate estimation of groundwater based on radioactive tracers (3H et 14C) in an unconfined aquifer on a semi-arid area, Iullemede Basin. *Niger J Hydrol* 254:145–156
- Samanta G, Clifford DA (2005) Preservation of inorganic arsenic species in groundwater. *Environ Sci Technol* 39(22):8877–8882. <https://doi.org/10.1021/es051185i>
- Sangaré A, Driouch Y, Salvi S, Féménias O, Siebenaller L, Belkasm M, Béziat D, Dahire M, Ntarmouchant A, Adil S, Debat P (2014) Géologie des minéralisations aurifères du gisement tardi-ébournéen de Kalana (Birimine, Sud-Ouest du Mali). *Bull L'institut Sci Rabat Section Sci Terre* 2014(36):85–108
- Smedley PL (1996) Arsenic in rural groundwater in Ghana: part special issue: Hydrogeochemical studies in sub-saharan Africa. *J Afr Earth Sci* 22:459–470

- Smedley PL, Knudsen J, Maiga D (2007) Arsenic in groundwater from mineralized proterozoic basement rocks of Burkina Faso. *Appl Geochem* 22:1074–1092
- Smedley L, Kinniburgh DG (2002) A review of the source, behavior and distribution of arsenic in natural waters. *Appl Geochem* 17(5):517–568
- Soumaila A, Konate M (2005) Caractérisation de la déformation dans la ceinture birimienne (paléoprotérozoïque) de Diagorou-Darbani (Liptako nigérien, Afrique de l’Ouest). *Afr Geo Rev* 13(3):161–178
- Soumaila A, Henry P, Rossy M (2004) Contexte de mise en place des roches basiques de la ceinture de roches vertes birimienne de Diagorou-Darbani (Liptako, Niger, Afrique de l’Ouest): plateau océanique ou environnement d’arc/bassin arrière-arc océanique. *C. R. Géoscience* 336:1137–1147
- Soumaila A (2000) Etude structurale, pétrographique et géochimique de la ceinture de Diagorou-Darbani, Liptako, Niger Occidental (Afrique de l’Ouest). Thèse, Univ. Franche- Comté, pp. 56–203.
- Tankari DA, Tidjani DA, Idder GT, Yadjil DL, N., Matsallabi A., Ambouta J. M. K., Feidt C., Sterckeman T., Echevarria G. (2014) Diagnostic de la contamination des eaux par les éléments traces métalliques dans la zone aurifère de Komabangou – Tillabéri. *Niger Int J Biol Chem Sci* 8(6):2849–2857
- Tankari AD, Ibrahim OZ, Guéro Y, Morel JL, Feidt C, Echevarria G (2019) Impacts of artisanal gold mining on soil, water and plant contamination by trace elements at Komabangou, Western Niger. *J Geochem Explor* 205(2019):106328
- Taylor S, McLennan S (1995) The geochemical evolution of the continental crust. *Rev Geophys* 33(2):241–265
- Viers J, Dupré B, Gaillardet J (2009) Chemical composition of suspended sediments in World Rivers: new insights from a new database. *Sci Total Environ* 407:853–868
- WHO (2017) Guidelines for drinking water quality, 4th edn. World Health Organization, Geneva