



HAL
open science

Brain hemodynamic and electrical activity during functional upper limb movements in chronic stroke

Camille Muller, Makii Muthalib, Gérard Dray, Binbin Xu, Denis Mottet, Jérôme Froger, Marion Delorme, S. Perrey, Isabelle Laffont, Karima Bakhti

► **To cite this version:**

Camille Muller, Makii Muthalib, Gérard Dray, Binbin Xu, Denis Mottet, et al.. Brain hemodynamic and electrical activity during functional upper limb movements in chronic stroke. IS-PRM/ESPRM/SPMFR 2022 Congress, Jul 2022, Lisbonne, Portugal. , 2022. hal-03887363

HAL Id: hal-03887363

<https://imt-mines-ales.hal.science/hal-03887363>

Submitted on 11 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

C. O. Muller^{1,2}, M. Muthalib^{1,2}, G. Dray¹, B. Xu¹, D. Mottet¹, J. Froger^{1,3}, M. Delorme^{1,3}, S. Perrey¹, I. Laffont^{1,2}, K.K.A. Bakhti^{1,2}

1 - EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Alès, Montpellier, France
 2 - Physical Rehabilitation and Medicine, CHU Montpellier, Montpellier, France
 3 - Physical Rehabilitation and Medicine, CHU Nîmes, Le Grau Du Roi, France

✉ camille.muller@umontpellier.fr

INTRODUCTION

Two-thirds of stroke survivors suffer from **persistent functional limitations** of their paretic upper limbs (UL). To cope with this persistent sequel, a good understanding of UL recovery mechanisms is needed. Portable functional neuro-imagery could be easily used for **brain-movement evaluation** to highlight **neuroplasticity mechanisms during rehabilitation**.

Objective: To fuse fNIRS and fEEG methods to evaluate sensorimotor cortical (SMC) activity during UL movements

METHODS

Participants:

- 3 chronic stroke patients (mean age = 59 years, >6 months post-stroke)
- Upper limb impairment (mean FM-UE = 48/66)

Experimental protocol (Fig.1):

- 3 blocs of 20s of each functional task (Fig. 1) with paretic and non-paretic UL
- Synchronized recording of brain (fNIRS, fEEG, Fig. 2) and movement (Kinect, graphic tablet pad) parameters




Fig 1. a) Reaching and b) Circular steering tasks

Fig 2. fNIRS and fEEG layout over the SMC region

DISCUSSION

• **Neuromonitoring** platform to explore movement-related neurovascular coupling mechanisms

• **Relative ease of installation** provides a **routine** way to explore level of neuroplasticity during stroke rehabilitation

• Long term objective: **precision medicine**

RESULTS




Fig 3. Movements of the paretic arm (a) and non-paretic arm (b) for the circular steering task in one patient

Kinematics (Fig. 3): lower performances with paretic UL




Fig 4. fNIRS hemodynamic response during circular movements

fNIRS provides (Fig. 4) information about:

- Level of SMC activation (**↑ oxygenated blood**)
- **Ipsilateral and contralateral SMC** difference




Fig 5. Power modification in Alpha and Beta bands during circular movements

fEEG provides (Fig. 5) information about:

- Level of **SMC excitability** (**↓ Alpha (8-13Hz) and Beta (14-35Hz) power**)
- **Ipsilateral and contralateral SMC** difference