



Virus-based biological control facing climate change

Miguel Lopez-Ferber

► To cite this version:

Miguel Lopez-Ferber. Virus-based biological control facing climate change. 18th meeting IOBC/WPRS Working Group Microbial and Nematode Control of Invertebrate Pests, Jun 2022, České Budějovice, Czech Republic. pp.109-114. hal-04169533

HAL Id: hal-04169533

<https://imt-mines-ales.hal.science/hal-04169533>

Submitted on 15 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Virus-based biological control facing climate change

Miguel Lopez-Ferber¹

¹*Hydrosiences Montpellier, Univ Montpellier, IMT Mines Ales, IRD, CNRS, 6 Avenue de Clavières, Alès, France, ²INRAE, SPE., Rte des Chappes, 06560 Sophia Antipolis, France*
Correspondence: miguel.lopez-ferber@mines-ales.fr

Abstract: Climate change will probably induce changes in the use of virus-based biocontrol agents at various levels. Climatic changes will affect host development, indirectly modifying virus behaviour. Invasive species will constitute potential new niches for virus populations, but also may reduce their traditional hosts. Introduction of new viruses adapted to such invasive species in their new geographic areas might be an appropriate strategy. Variation in UV radiation, in rain or wind will directly affect the survival or the availability of the virus for the appropriate hosts. Variation in temperatures influence insect host development, but also plant development, and will indirectly affect virus/host relationships.

Instructions of use, timing and doses should be re-evaluated in function of new climate parameters. In addition, formulation changes might be required to optimise the persistence of virus-based products in new environmental conditions.

Key words: pests, diseases, integrated control, climatic change, adaptation

Introduction

Climate change is now well established and is affecting us in many different ways. Abundant literature has been devoted to the effects of climate changes on agriculture, mainly concentrating on the ability of crops to cope with increasing temperatures, or with drought, and the ways to avoid production losses, by changing varieties or crop species by others more adapted to the new climatic conditions. These adaptations often imply changes in the pest affecting the crops, and thus, in the way to control them.

Another consequence of the climate change is the colonisation of new areas by invasive species, that might acquire the pest status in the new area. Fighting these invasive species may require the adaptation of native pathogens or predators. Alternatively, a solution could be the introduction of the agents that regulate the invader in its original distribution area, where often it does not represent a problem.

In temperate regions, increasing the temperature will accelerate the development of insects, leading to a higher generation number in multivoltine insect species in temperate regions. Similarly, increasing the period of appropriate temperature will have similar consequences. This effect will of course have an upper limit that is the maximal temperature supported by the insect.

However, the most limiting factor for larvae survival seems to be the lowest temperature that larvae can stand during diapause. The work by Causse (1976) showed that the density of first generation is clearly influenced by the lowest temperature during the winter. Climatic changes, by inducing higher extreme might also have this effect.

Whatever situation, climatic change will affect the life cycle of the insect host, and thus, that of the virus pathogen. From an agronomic perspective, that probably means adapting the treatments, maybe having more treatment per season.

When addressing specifically the virus based biocontrol agents, all these concerns should be taken into account. In addition, there are other points to consider, due to the specificity of the virus. In this work we will address more specifically the insect pathogenic viruses that are or could be used to control insect pests. Even if at the present time baculovirus are almost the only virus group used for as biocontrol agents, I'll try to depict a more general view, that could be applied to other virus families.

Invasive insect species: adaptation of virus populations

Adaptation of a virus to a new host species has been described both in laboratory conditions and in fields. Determination of baculovirus host range determinants was a hot research topic in the 90's. Using the model baculoviruses AcMNPV and BmNPV, Maeda's group in Japan and Croizier's group in France demonstrate that AcMNPV could be adapted to replicate in silkworm and silkworm cells by modifying the virus helicase gene (Kondo and Maeda, 1991; Crozier et al., 1994; Argaud et al., 1998). Other works revealed that there was often possible to adapt(baculo)viruses to new hosts, but that the genes involved are not allways the same, preventing a systematic approach (Thiem and Cheng, 2009; Williams et al., 2022).

Field observations confirm the ability of a local virus population to adapt to a new host. The guatemala tuber moth, *Tecia solanivora* progressively invaded south America and recently was found in the Canary Islands (Puillandre et al., 2008). This pest displaced the potato tuber moth, *Phthorimaea operculella*, but the two species coexist in many locations. The *Phthorimaea operculella* granulovirus is found associated to all *P. operculella* populations. The isolates of this virus collected were or when no *T. solanivora* is present do not replicate in this species. However, soon after invasion, granuloviruses able to replicate in *T. solanivora* larvae were isolated in Venezuela (Niño de Gualdron and Notz, 2000). The process of invasion of *T. solanivora* in Colombia has been analysed in the 2000's (Zeddarn et al., 2013). Sampling for viruses was carried out and revealed that the genetic composition of virus isolates changed to adapt to the presence of this new potential host (and probably to the reduction of the original host). This adaptation process will probably take place for every invasion, although the success cannot be granted.

A complementary or alternative way is the introduction of a virus already adapted to this species. This approach has often been taken both for invasive species or for already established pests. As example, the mexican isolate of *Cydia pomonella* granulovirus (CpGV-M), the first discovered, was used for developing virus based insecticides to control codling moth in european orchards.

Sensitivity to environmental conditions

The primary infection is a key point that may be heavily affected by climatic change. Transmission will not be affected for viruses that are propagated by close contact of individuals. This can be the situation for social insects living in colonies (bees, termites, ...), gregarious insects, or those in high density situation. A well known example is *Oryctes* virus used for the control of the rhinoceros beetle, *Oryctes rhinoceros* (Huger, 2005). Two main ways of contamination have been proposed. Virus replicate actively in the midgut epithelium, and particles are released in high numbers on the feces, contaminating feeding burrows of palm crows. The *Oryctes* virus is a non occluded virus, whose particle and replication remembers that of baculoviruses. It was classified as a "non occluded baculovirus" before an independent family was proposed in the 2000's. This virus is highly susceptible to environmental

conditions, heat and UV radiation, but the huge amounts released and the shield offered by palm leaf certainly attenuate such limitations. In addition, the second way of transmission of the infection, that is supposed to be the most important, is by direct contact during mating or cooccupation of the same palm. In such circumstances, the delay on the external environment is very reduced. The same constraints will apply to other viruses with similar transmission strategies.

Viruses that need to survive into the environment between infection cycles will be affected by changes in climate. If the virus is supposed to survive in the environment between infections, variation in the climatic conditions may affect its persistence. Increasing temperature is probably not a direct limiting factor for survival of these viruses. Baculovirus OBs can stand temperatures up to 50 °C without significantly losing infectivity (McLeod et al., 1977). The same applies for other occluded insect viruses, like cytoplasmic polyhedrosis viruses (Reoviridae) (Tanada and Chang, 1968). Baculovirus can survive as OBs to low temperatures without significantly losing pathogenicity. Accordingly, stronger or milder winter temperatures should not greatly influence the virus survival. However, when transmission between years is mainly dependent on a diapausing stage, like in vertical transmission, or in chronically infecting viruses, the survival of the insect may be affected, and indirectly, that of the virus. In addition, increasing the average temperatures will induce faster insect development, and thus an increasing number of generations per year. From an agronomic perspective, this means expanding the treatment period.

Increasing the sun light irradiance implies a higher UV dosis, that affects virus survival. Increasing rainfall intensity may result on virus wash off from the aerial parts of the plant. Similarly, strong winds may also affect the disponibility of the virus to the insect pest. Baculovirus inactivation by UV light has been studied both in laboratory and in field conditions. It is one of the key point that limit the development of this crop protection biocontrol agents in the actual climatic conditions. To increase the survival time, commercial products include UV protection, like flurescent brigtners or other UV protective substances (Çakmak et al., 2021). If the changes on UV radiation are significant, it may be necessary to reevaluate the level of UV protection required, by increasing the concentration of UV protector, or by changing the substance used.

Brassel and Bentz (1979) reported a 5.6 increase in the resistance to UV inactivation on CPGV after selection. In their work they do not specify which isolate they used, but it was probably the CpGV-M, that is know to contain extremely low genetic diversity, which may explain the relatively low response to the selection process. Similar approaches were taken for other baculoviruses, confirming a genetic variability for the resistance to UV light, that could be exploited when developping biocontrol agents, and that the virus could mobilize for its adaptation to evolving climatic conditions.

In *Chrysodeixis chalcites* nucleopolyhedrovirus, two genes homologous to photolyases were found by when sequencing the entire virus genome, *phr1* and *phr2*. The DNA repair activity was demonstrated for *phr2*, but not for *phr1* (van Oers et al., 2008). However, homology searches have detected homologous genes in group II alphabaculoviruses infecting plusiine insects (Xu, 2008), but not in other baculoviruses.

Overexpression of the Bm65 virus gene results in a higher UV tolerance. Bm65, homologous to Ac79, is predicted to be an endonuclease, that could be involved in UV-induced DNA repair (Tang et al., 2015). Tang et al. (2019) observed that BmN cells infected with BmNPV show higher resistance to high UV radiation than non infected cells, suggesting than the baculovirus activate UV reparing pathways, either by producing UV repair enzymes or by complementing cellular pathways.

Other virus families also encode photolyases. The activity of the *Amsacta moorei* entomopoxvirus gene 25 (AMV025) is able to rescue photolyase deficient *E. coli* cells, confirming its activity as a functional photolyase (Nalcacioglu et al., 2010).

A recent paper by a sudafrikan team (Mwanza et al., 2022), describes selection for UV resistance on a betabaculovirus local isolate, the *Cryptophlebia Leucotreta* Granulovirus-SA. These authors obtain a more than 1000 fold improved pathogenicity after only five passages on insects after UV exposure. The authors did not conclude on the genes involved in UV tolerance or resistance, but address the point that UV resistance probably has a fitness cost, as it is not generally distributed in the virus population.

As increasing the resistance to UV radiation in baculovirus is a key point in the development of biocontrol agents, research was done by inserting DNA repair enzymes from other viruses (Petrick et al., 2003). These authors did not obtain a significant decrease in the inactivation rate for OBs.

Increasing rainfall or wind intensity will wash off the virus from the leaves, but may also mobilize the virus present in the soil. Most probably, the adhesion of the virus particle to a surface is a pasive mechanism. Crude virus preparations of *Heliothis zea* nucleopolyhedrovirus strongly adhere to surfaces (Ignoffo et al., 1997), probably due to the soluble components of the dead larva tissues, but this does not seem to be a general feature. Virus-based insecticides can be supplemented with adjuvants that increase the adhesion of particles to the surfaces. Soil is a reservoir for virus, protecting it from UV. If the target passes through soil, it can be contaminated. Alternatively, heavy rain or wind can transport virus from the soil to the aerial parts of plants (Jacques, 1985).

Tritrophic interactions

Andrews and Sikorovski (1973), cite in Jacques, 1985, noted that deposits of HNPV on cotton leaves were inactivated during the night when dew caused the deposits to be wet. Plants secrete substances that degrade the virus, maybe combined with humidity. Secretion is a response to stress, that can be climatic (higher temperatures, lower water availability), and accordingly, it might be necessary to adjust formulations to varying exterior temperatures. A more subtle interaction is the response of plants by making the upper parts of the leaves thicker, and sclerotised, and thus, not eatable to larvae. If the virus is located on this upper part, the probability of ingestion will be lower.

Conclusion

Climatic change will affect the use of viruses as biocontrol agents for the regulation of insect pests. It will offer opportunities to develop new approaches, but direct transposition from what we know to work in one place could not be done to new locations. A careful analysis of the local climate will be necessary. If introduction of a imported virus is the preferred solution, the similarity of climatic conditions between the virus original region and the intended location should be taken into account.

References

- Argaud, O., Croizier, L., Lopez-Ferber, M. and Croizier, G. 1998. Two key mutations in the host-range specificity domain of the p143 gene of *Autographa californica* nucleopolyhedrovirus are required to kill *Bombyx mori* larvae. J. Gen. Virol. 79 (Pt. 4): 931-935.

- Brassel, J.; Benz, G. 1979. Selection of a strain of the granulosis virus of the codling moth with improved resistance against artificial ultraviolet radiation and sunlight. *J. Invertebr. Pathol.* 33: 358-363.
- Çakmak, T., Simón, O., Kaydan, M. B., Tange, D. A., González Rodríguez, A. M., Piedra-Buena Díaz, A., Caballero, P., Hernandez Suarez, E. 2021. Effects of several UV-protective substances on the persistence of the insecticidal activity of the Alphabaculovirus of *Chrysodeixis chalcites* (ChchNPV-TF1) on banana (*Musa acuminata*, Musaceae, Colla) under laboratory and open-field conditions. *PLoS ONE* 16(5): e0250217. <https://doi.org/10.1371/journal.pone.0250217>.
- Cause, R. 1976. Etude de la localisation et de la mortalité hivernale des larves de *Laspeyresia pomonella* L. (*Lepidoptera, Tortricidae*) en vergers modernes de pommiers de la basse vallée du Rhône. *Ann. Zool. Ecol. Animal* 8 (1): 83-101.
- Croizier, G., Croizier, L., Argaud, O., Poudevigne, D. 1994. Extension of *Autographa californica* nuclear polyhedrosisvirus host-range by interspecific replacement of a short DNA-sequence in the P143 helicase gene. *Proc. Natl. Acad. Sci. USA* 91(1): 48-52.
- Huger, A. M. 2005. The Oryctes virus: Its detection, identification, and implementation in biological control of the coconut palm rhinoceros beetle, *Oryctes rhinoceros* (Coleoptera: Scarabaeidae). *J. Invertebr. Pathol.* 89: 78-84.
- Ignoffo, C. M., Garcia, C., Saathoff, S. G. 1997. Sunlight Stability and Rain-Fastness of Formulations of Baculovirus heliothis. *Environmental entomology* 22(6) 1470-1474.
- Jacques, R. 1985. Stability of entomopathogenic viruses. *Misc. publ. Entomol. Soc. Am.* 10 (3): 99-116.
- Kondo, A. and Maeda, S. 1991. Host range expansion by recombination of the baculoviruses *Bombyx mori* nuclear polyhedrosis virus and *Autographa californica* nuclear polyhedrosis virus. *J. Virol.* 65: 3625-3632.
- McLeod, P. J., Yearian, W. C., Young III, S. Y. 1977. Inactivation of *Baculovirus heliothis* by ultraviolet irradiation, dew, and temperature. *J. invertebr. Pathol* 30(2): 237-241.
- Mwanza, P., Jukes, M., Dealtry, G., Lee, M. Moore, S. 2022. Selection for and Analysis of UV-Resistant *Cryptophlebia leucotreta* Granulovirus-SA as a Biopesticide for *Thaumatotibia leucotreta* Viruses 10.3390/v14010028.
- Nalcacioglu, R., Dizman, Y. A., Vlak, J. M., Demirbag, Z., van Oers, M. M. 2010. Amsacta moorei entomopoxvirus encodes a functional DNA photolyase (AMV025) *J. Invertebr. Pathol.* 105(3): 363-365. doi: 10.1016/j.jip.2010.06.013.
- Niño de Gualdrón, L. and Notz, A. 2000. Patogenicidad de un virus granulosis de la polilla de la papa *Tecia solanivora* (Povolny) 1973. (*Lepidoptera: Gelechiidae*), en el estado Mérida, Venezuela. *Boletín Entomologia Venezolana* 15(1): 39-48.
- Petrik, D. T., Angela Iseli, A., Montelone, B. A., Van Etten, J. L., Clem, R. J. 2003: Improving baculovirus resistance to UV inactivation: increased virulence resulting from expression of a DNA repair enzyme. *J. Invertebr. Pathol.* 82: 50-56
- Puillandre, N., Dupas, S., Dangles, O., Zeddami, J.-L., Capdevielle-Dulac, C., Barbin, K., Torres-Leguizamon, M., Silvain, J.-F. 2008. Genetic bottleneck in invasive species: the potato tuber moth adds to the list. *Biol. Invasions* 10: 319-333. DOI 10.1007/s10530-007-9132-y-
- Tanada, Y., and Chang, G. Y. 1968- Resistance of the alfalfa caterpillar, *Colias eurytheme*, at high temperatures to a cytoplasmic-polyhedrosis virus and thermal inactivation point of the virus. *J. Invertebr. Pathol.* 10(1): 79-83.
- Tang, Q., Hu, Z., Yang, Y., Wu, H., Qiu, L., Chen, K., Li, G. 2015. Overexpression of Bm65 correlates with reduced susceptibility to inactivation by UV light. *J. Invertebrate Pathol.* 127: 87-92.

- Tang, Q., Chen, F., Wu, P., Qiu, L., Chen, H., Chen, K., Li, G. 2019. BmNPV infection correlates with the enhancement of the resistance of *Bombyx mori* cells to UV radiation. Arch. Insect Biochem. Physiol. 102(1): e21598. DOI 10.1002/arch.21598.
- Thiem, S. M., Cheng, X.-W. 2009. Baculovirus host range. Virologica Sinica 24(5): 436-457.
- van Oers, M. M., Lampen, M. H., Bajek, M. I., Vlak, J. M., Eker, A. P. M. 2008. Active DNA photolyase encoded by a baculovirus from the insect *Chrysodeixis chalcites*. DNA Repair 7(8): 1309-1318.
- Williams, T., Lopez-Ferber, M., Caballero, P. 2022. Nucleopolyhedrovirus Coocclusion Technology: A New Concept in the Development of Biological Insecticides. Front. Microbiol. 12: 810026. Doi: 10.3389/fmicb.2021.810026.
- Xu, F., Vlak, J. M., van Oers, M. M. 2008. Conservation of DNA photolyase genes in group II nucleopolyhedroviruses infecting plusiine insects. Virus Research 136(1-2): 58-64. doi: 10.1016/j.virusres.2008.04.017.
- Zeddai, J.-L., Léry, X., Gómez-Bonilla, Y., Espinel-Correal, C., Páez, D., Rebaudo, F., López-Ferber, M. 2013. Responses of different geographic populations of two potato tuber moth species to genetic variants of *Phthorimaea operculella* granulovirus. Entomol. Exp. Appl. 149(2): 138-147. Doi: 10.1111/eea.12115.