



HAL
open science

Letter to the Editor: Response regarding “Single-session anodal transcranial direct current stimulation to enhance sport-specific performance in athletes: A systematic review and meta-analysis”

Tom Maudrich, Patrick Ragert, S. Perrey, Rouven Kenville

► To cite this version:

Tom Maudrich, Patrick Ragert, S. Perrey, Rouven Kenville. Letter to the Editor: Response regarding “Single-session anodal transcranial direct current stimulation to enhance sport-specific performance in athletes: A systematic review and meta-analysis”. *Brain Stimulation*, 2023, 16 (5), pp.1551-1552. 10.1016/j.brs.2023.09.001 . hal-04442115

HAL Id: hal-04442115

<https://imt-mines-ales.hal.science/hal-04442115>

Submitted on 6 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Letter to the Editor: Response regarding “Single-session anodal transcranial direct current stimulation to enhance sport-specific performance in athletes: A systematic review and meta-analysis”

Dear Editor,

We appreciate the valuable input provided by Machado & Amiri (2023). Through this response letter, our aim is to eliminate any uncertainties pertaining to our meta-analysis [1]. Please note that substantive remarks by the authors are condensed for the sake of readability.

An initial comment concerned the inclusion of 1) tDCS studies using multiple target areas and 2) tDCS studies combining stimulation with interventions. Regarding the first point [2], used a dual montage covering both temporal cortex (TC) and primary motor cortex (M1). Crucially, both targets were referenced separately. Such montages are employed to dissect distinct effects arising from two specific target areas within interventions utilizing dual-stimulation protocols. This approach gains particular importance when discrete ergogenic outcomes are associated with each target area, similar to the context highlighted in the study of [2]. When a common reference is used to cover both target areas, the challenge of drawing area-specific inferences becomes intricate due to the dominant influence of the common reference. This can result in potential neural interference or cross-talk, complicating the accurate interpretation of findings. Consequently, the inclusion of [2] was undertaken to address this concern. It acknowledges that using separate references for each target area, as opposed to a common reference, potentially provides a more robust basis for deriving definitive conclusions about the distinct influences exerted by these individual target areas. In another study [3], both M1 hand areas received stimulation, while reference electrodes were placed on spinous processes. This electrode arrangement creates a distinct current flow for this montage, although the main focus of stimulation is on M1. Furthermore, it is important to note that the positioning of reference electrodes is a known factor affecting the variability of tDCS protocols, along with other aspects like current intensity, electrode size, current density, and stimulation duration. For example, a reference electrode using conventional tDCS setups is not necessarily ineffective in modulating underlying brain activity [4]. Therefore, tDCS always modulates several brain regions simultaneously. If we were to overly standardize all these factors, it would make it difficult to combine analyses, identify patterns, or conduct follow-up studies based on them. We are, therefore, confident that the standardization of inclusion criteria adopted in our meta-analysis is restrictive enough to include only studies that are closely related in terms of content and methodology, yet permissive enough to account for the heterogeneity of tDCS protocols. Crucially, all these implications were transparently mentioned as well as discussed in

our original meta-analysis.

Concerning 2) [5–7], used a type of Stroop task to induce mental fatigue. Both [5,6] applied tDCS (a-tDCS or sham) during the mental fatigue task. Nikoohaarf Salehi et al. [7] induced mental fatigue before tDCS was applied. In all studies, sport-specific performance was assessed after completion of tDCS. We agree that the induction of mental fatigue in addition to brain stimulation might have influenced the outcome of tDCS. However, we opted to include these studies, since athletes experience mental fatigue and cognitive challenges such as anxiety and performance pressure before competitions, making these studies important for the overall aim of the meta-analysis to investigate the ecological validity of tDCS, or as we explained “investigation of sport-specific performance changes through tDCS seems essential to further approach the understanding of the potential of tDCS applications under real-life conditions compared to highly controlled laboratory settings”.

Another point was raised concerning the inclusion of a study by Fortes et al. [8]. Machado & Amiri (2023) assumed that participants should not be eligible for inclusion because they were recreational practitioners, citing Fortes et al. [8] as follows: “Twelve participants were recreationally trained in resistance training”. Crucially, the complete statement of the article in question reads: “Twelve participants were recreationally trained in resistance training (i.e., individuals consistently trained from three to ten years, frequency 3–5 sessions per week)”. Accordingly, the trained subgroup certainly fits within our athlete inclusion criteria as we included “healthy male or female adult athletes (participating regularly in organized sport for at least 2 years before the experiment)”. Importantly, only data from trained participants of this study was included in our meta-analysis.

Moreover, Machado & Amiri (2023) raised concern regarding a study by da Silva Machado et al. [9], where HD-tDCS and conventional tDCS were employed. This study was included in our original meta-analysis, although we chose to include only the data of the conventional tDCS setup of their study for the following reasons: 1) the majority of studies included in the meta-analysis used conventional tDCS, 2) the same brain target (M1) was stimulated, using the same participant sample, so the inclusion of both study arms may have influenced the overall effect. However, to check whether inclusion of the HD-tDCS arm in our meta-analysis would have changed the overall result, we conducted a reanalysis. The overall effect favouring a-tDCS over sham remained the same (SMD = 0.31, $p = 0.0002$), while the subgroup effect for M1 stimulation changed from SMD = 0.33 ($p = 0.01$) to SMD = 0.32 ($p = 0.008$). Therefore, we are confident in our initial

DOI of original article: <https://doi.org/10.1016/j.brs.2022.11.007>.

<https://doi.org/10.1016/j.brs.2023.09.001>

Received 30 August 2023; Accepted 2 September 2023

Available online 30 October 2023

1935-861X/© 2023 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

result. Furthermore, the authors seemed concerned with a subsequent statement in our original manuscript that reads: “One study compared conventional tDCS with HD-tDCS [9]. Because there were no differences between conventional tDCS and HD-tDCS in this study, the focus of the current review was on conventional tDCS.” This wording might have led the authors to believe we generally excluded HD-tDCS, which was not the case, as this statement referred exclusively to the study of da Silva Machado et al. [9].

A final issue concerned the practical implications resulting from a combination of findings from both subgroup analyses. We agree, that such implications cannot be drawn, and want to stress that we explicitly state that separate subgroup analyses were performed. At no point in the original article do we draw conclusions based on a combination of both subgroup analyses.

We hope this letter resolves the ambiguities cited and encourage future studies on this vibrant, dynamic topic of tDCS-induced performance enhancement.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was supported by the Max-Planck-Society.

References

- [1] Maudrich T, Ragert P, Perrey S, Kenville R. Single-session anodal transcranial direct current stimulation to enhance sport-specific performance in athletes: a systematic review and meta-analysis. *Brain Stimul* 2022;15(6):1517–29.
- [2] Kamali A-M, Saadi ZK, Yahyavi S-S, Zarifkar A, Aligholi H, Nami M. Transcranial direct current stimulation to enhance athletic performance outcome in experienced bodybuilders. *PLoS One* 2019;14(8).
- [3] Kamali AM, Kazemiha M, Keshtkarhesamabadi B, Daneshvari M, Zarifkar A, Chakrabarti P, et al. Simultaneous transcranial and transcutaneous spinal direct current stimulation to enhance athletic performance outcome in experienced boxers. *Sci Rep* 2021;11(1):19722.
- [4] Nitsche MA, Doemkes S, Karakose T, Antal A, Liebetanz D, Lang N, et al. Shaping the effects of transcranial direct current stimulation of the human motor cortex. *J Neurophysiol* 2007;97(4):3109–17.
- [5] Penna EM, Edson, Campos BT, Ferreira RM, Parma JO, Lage GM, et al. No effects of mental fatigue and cerebral stimulation on physical performance of master swimmers. *Front Psychol* 2021;12.
- [6] Fortes LS, Faro H, de Lima-Junior D, Albuquerque MR, Ferreira MEC. Non-invasive brain stimulation over the orbital prefrontal cortex maintains endurance performance in mentally fatigued swimmers. *Physiol Behav* 2022;250:113783.
- [7] Nikooharf Salehi E, Jaydari Fard S, Jaberzadeh S, Zoghi M. Transcranial direct current stimulation reduces the negative impact of mental fatigue on swimming performance. *J Mot Behav* 2022;54(3):327–36.
- [8] Fortes LS, Mazini-Filho M, Lima-Júnior D, Machado DGS, Albuquerque MR, Fonseca FS, et al. Transcranial stimulation improves volume and perceived exertion but does not change power. *Int J Sports Med* 2021;42(7):630–7.
- [9] da Silva Machado DG, Bikson M, Datta A, Caparelli-Dáquer E, Unal G, Baptista AF, et al. Acute effect of high-definition and conventional tDCS on exercise performance and psychophysiological responses in endurance athletes: a randomized controlled trial. *Sci Rep* 2021;11(1):13911.

Tom Maudrich^{*}, Patrick Ragert

Department of Movement Neuroscience, Faculty of Sport Science, Leipzig University, Leipzig, Germany

Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

Stéphane Perrey

EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Ales, Montpellier, France

Rouven Kenville

Department of Movement Neuroscience, Faculty of Sport Science, Leipzig University, Leipzig, Germany

Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

^{*} Corresponding author.

E-mail address: tom.maudrich@uni-leipzig.de (T. Maudrich).

Handling Editor: Dr. M. George